



CLINICAL PRACTICE GUIDELINES

EMT

**EMERGENCY
MEDICAL TECHNICIAN
- incorporating BTEC**

2021 Edition



PHECC Clinical Practice Guidelines

First Edition, 2001

Second Edition, 2004

Third Edition, 2009

Third Edition, Version 2, 2011

Fourth Edition, April 2012

Fifth Edition, July 2014

Sixth Edition, March 2017 (Updated February 2018)

Seventh Edition, August v2 2021

Published by:

Pre-Hospital Emergency Care Council

2nd Floor, Beech House, Millennium Park, Osberstown, Naas, Co Kildare, W91 TK7N, Ireland.

Phone: +353 (0)45 882042

Email: info@phecc.ie

Web: www.phecc.ie

ISBN 978-0-9955288-8-8

© Pre-Hospital Emergency Care Council 2021

Permission is hereby granted to redistribute this document, in whole or part, for educational, non-commercial purposes providing that the content is not altered and that the Pre-Hospital Emergency Care Council (PHECC) is appropriately credited for the work. Written permission from PHECC is required for all other uses. Please contact the author: r.carney@phecc.ie

Table of Contents

EMERGENCY
MEDICAL TECHNICIAN
- incorporating BTEC

FOREWORD	4
ACCEPTED ABBREVIATIONS	5
ACKNOWLEDGEMENTS	8
INTRODUCTION	10
IMPLEMENTATION AND USE OF CLINICAL PRACTICE GUIDELINES	11

CLINICAL PRACTICE GUIDELINES

INDEX	13
KEY/CODES EXPLANATION	15
SECTION 1 Principles of general care	16
SECTION 2 Airway and Breathing	22
SECTION 3 Cardiac	28
SECTION 4 Circulation	30
SECTION 5 Medical	31
SECTION 6 Neurological	34
SECTION 7 Behavioural and Mental Health Emergencies	38
SECTION 8 Trauma	40
SECTION 9 Environmental	47
SECTION 10 Toxicology	49
SECTION 11 Infectious	51
SECTION 12 Maternal.....	52
SECTION 13 Paediatric	58
SECTION 14 Resuscitation	81
SECTION 15 Palliative Care	88
SECTION 16 Operations	89
APPENDIX 1 Medication Formulary	92
APPENDIX 2 Medications & Skills Matrix	114
APPENDIX 3 Critical Incident Stress Management (CISM)	123
APPENDIX 4 CPG Updates for Emergency Medical Technician	126

This Handbook comprises the 2021 Edition Clinical Practice Guidelines (CPGs). These guidelines outline patient assessments and pre-hospital management for responders at:

RESPONDER LEVEL

- Cardiac First Responder
- First Aid Responder
- Emergency First Responder

REGISTERED PRACTITIONER

- Emergency Medical Technician
- Paramedic
- Advanced Paramedic



I am delighted that there are now 357 CPGs in total to guide integrated care across the six pre-hospital emergency care clinical levels. These CPGs ensure that responders and practitioners are practicing to best international standards and support PHECC's vision that people in Ireland receive excellent pre-hospital emergency care.

I would like to acknowledge the hard work and commitment the members of the Medical Advisory Committee have shown during the development of this publication, guided by Dr David Menzies (Chair). A special word of thanks goes to Dr Brian Power who retired in 2020 and has made an enormous contribution to the advancement of pre-hospital emergency care in Ireland. I want to acknowledge the PHECC Executive, for their continued support in researching and compiling these CPGs and paving the way for the future development of the pre-hospital emergency care continuum.

I recognise the contribution made by many responders and practitioners, whose feedback has assisted PHECC in the continual improvement and development of CPGs and welcome these guidelines as an important contribution to best practice in pre-hospital emergency care.

A handwritten signature in black ink that reads "Jacqueline Burke".

Dr Jacqueline Burke, Chairperson
Pre-Hospital Emergency Care Council

Advanced Paramedic	AP
Advanced Life Support	ALS
Airway, Breathing & Circulation	ABC
All Terrain Vehicle	ATV
Altered Level of Consciousness	ALoC
Automated External Defibrillator	AED
Bag Valve Mask	BVM
Basic Life Support	BLS
Blood Glucose	BG
Blood Pressure	BP
Basic Tactical Emergency Care	BTEC
Capillary Refill Time	CRT
Carbon Dioxide	CO ₂
Cardiopulmonary Resuscitation	CPR
Cervical Spine	C-spine
Chronic Obstructive Pulmonary Disease	COPD
Clinical Practice Guideline.....	CPG
Continuous Positive Airway Pressure.....	CPAP
Degree.....	°
Degrees Celsius.....	°C
Dextrose (Glucose) 10% in water.....	D ₁₀ W
Dextrose (Glucose) 5% in water	D ₅ W
Do Not Resuscitate.....	DNR
Drop (gutta)	gtt
Electrocardiogram	ECG
Emergency Department	ED
Emergency Medical Technician	EMT
Endotracheal Tube	ETT

Foreign Body Airway Obstruction	FBAO
Fracture	#
General Practitioner.....	GP
Glasgow Coma Scale	GCS
Gram.....	g
Intramuscular	IM
Intranasal	IN
Intraosseous	IO
Intravenous.....	IV
Joules	J
Kilogram	kg
Laryngeal Mask Airway	LMA
Mean Arterial Pressure	MAP
Medical Practitioner.....	MP
Microgram	mcg
Milligram.....	mg
Millilitre.....	mL
Millimole.....	mmol
Minute	min
Modified Early Warning Score.....	MEWS
Motor Vehicle Collision	MVC
Myocardial Infarction	MI
Milliequivalent	mEq
Millimetres of mercury	mmHg
Nasopharyngeal airway	NPA
Nebulised	NEB
Negative decadic logarithm of the H ⁺ ion concentration	pH

Orally (per os)	PO
Oropharyngeal airway.....	OPA
Oxygen	O ₂
Paramedic.....	P
Peak Expiratory Flow Rate.....	PEFR
Per rectum	PR
Per vagina	PV
Percutaneous Coronary Intervention	PCI
Personal Protective Equipment	PPE
Psychiatric Nurse	PN
Pulseless Electrical Activity	PEA
Pulseless Ventricular Tachycardia	pVT
Respiration rate	RR
Return of Spontaneous Circulation.....	ROSC
Revised Trauma Score	RTS
Saturation of arterial Oxygen	SpO ₂
Spinal Motion Restriction	SMR
ST Elevation Myocardial Infarction	STEMI
Subcutaneous.....	SC
Sublingual.....	SL
Supraventricular Tachycardia	SVT
Systolic Blood Pressure	SBP
Therefore	∴
Total body surface area	TBSA
Ventricular Fibrillation.....	VF
Ventricular Tachycardia	VT
When necessary (pro re nata)	prn

The process of developing CPGs has been long and detailed. The quality of the finished product is due to the painstaking work of many people, who through their expertise and review of the literature, ensured a world-class publication.

PROJECT LEADS

Mr Ricky Ellis, Programme Manager, PHECC

Mr Raymond Carney, Programme Manager, PHECC

MEDICAL ADVISORY COMMITTEE

Dr David Menzies (Chair), Consultant in Emergency Medicine, Member of Council

Mr David Irwin (Vice Chair), Advanced Paramedic, Nominated by Chair

Professor Gerard Bury, Director, UCD Centre for Emergency Medical Science

Mr Ian Brennan, Advanced Paramedic, Operational Resource Manager, HSE NAS

Mr Hillery Collins, Paramedic, Vice Chairperson of Council

Dr Niamh Collins, Consultant in Emergency Medicine, Connolly Hospital, Blanchardstown

Mr Eoghan Connolly, Advanced Paramedic, representative from the Irish College of Paramedics

Dr Lisa Cunningham Guthrie, Registrar in Emergency Medicine, Nominated by Chair

Mr Mark Dixon, Advanced Paramedic, Paramedic Programme Lead,
Graduate Entry Medical School, University of Limerick

Mr David Hennelly, Advanced Paramedic, Clinical Development Manager, HSE NAS

Mr Macartan Hughes, Chief Ambulance Officer – Education and Competency Assurance, HSE NAS

Dr Shane Knox, Assistant Chief Ambulance Officer - Education Manager, HSE NASC

Dr Stanley Koe, Consultant Paediatrician, CHI at Tallaght & Connolly Hospitals

Dr Mick Molloy, Consultant in Emergency Medicine, Wexford General Hospital

Mr Shane Mooney, Education and Competency Assurance Officer, HSE NAS

Dr Peter O'Connor, Medical Director, Dublin Fire Brigade

Professor Cathal O'Donnell, Clinical Director, HSE NAS

Mr Martin O'Reilly, District Officer, EMS Support, Dublin Fire Brigade

Dr Jason van der Velde, Clinical Lead, MEDICO Cork, Member of Council

Dr Philip Darcy, Consultant in Emergency Medicine, Connolly Hospital, Blanchardstown

EXTERNAL CONTRIBUTORS

PHECC would like to thank and acknowledge all of the experts who contributed to the creation of these Clinical Practice Guidelines.

SPECIAL THANKS

An extra special thanks to all the PHECC team who were involved in this project, especially Margaret Bracken, Aisling Ryan and Ashling Weldon for their painstaking recording of details and organisational skills.

MEDICATION FORMULARY REVIEW

Ms Regina Lee, MPSI

EXTERNAL CLINICAL REVIEW

Responder

Niamh O'Leary

Michelle O Toole

Emergency Medical Technician

Gareth Elbell

Gavin Hoey

Paramedic

Eithne Scully

Andy O Toole

Advanced Paramedic

Terry Dore

Pete Thorpe

EXTERNAL QUALITY REVIEW

Dr Jack Collins

Welcome to the 2021 edition of the PHECC Clinical Practice Guidelines. This edition has been a long time in development and reflects the significant effort and contribution to the new CPGs by so many people.

As ever, a robust development and review process has been applied to the new and revised CPGs, including a detailed and comprehensive quality assurance process.

Pre-Hospital Care in Ireland has evolved significantly since the first editions of the CPGs. The suite of care the CPGs now enable is progressive and transformative across all levels of responder and practitioner.

The impact of Covid-19 has influenced these CPGs, both in posing challenges in continuing the regular Medical Advisory Committee meetings and discussions while also giving rise to a specific suite of vaccination CPGs that enable PHECC practitioners to support the national Covid-19 vaccination programme.

For the first time, we have CPGs that enable practitioners to not convey patients to hospital as a matter of default. The non-conveyance CPGs are a step towards more alternative care pathways for our patients, in recognition that the traditional hospital-centric model for emergency care is not always appropriate or feasible. This suite of non-conveyance CPGs will be a key area for expansion and development in the next term of the Medical Advisory Committee.

Further developments include the designation of certain CPGs and elements of other CPGs as 'non-core'. This non-core element replaces the previous process of 'exemptions' accommodated for certain CPGs and recognises that not all Licenced CPG Providers need to implement every single CPG.

I would like to express my sincere thanks to all who contributed to this edition of the CPGs including the members of the Medical Advisory Committee, those who submitted queries for consideration, speciality groups and clinical programmes who provided expert external advice and feedback.

In particular, I would like to thank Dr Brian Power who retired from PHECC in 2020. Brian created the first edition of the PHECC CPGs and has managed the process of CPG development since then, including the majority of the development work for this suite of CPGs. Brian's contribution to the advancement of pre-hospital emergency care in Ireland has been significant and is the framework that supports responders and practitioners still. Since Brian's retirement, Ricky Ellis kindly and ably stepped into the gap, continuing to support MAC in the finalisation of the CPGs before handing over to Ray Carney, PHECC's new Clinical Programme Manager. Thank you both.

Finally, thanks to you, the responders and practitioners who implement these CPGs. I believe these CPGs will enable you to continue to provide expert compassionate pre-hospital care to patients every day of the year. PHECC greatly values your work and also your feedback.



Dr David Menzies, Chair Medical Advisory Committee



Clinical Practice Guidelines (CPGs) and the practitioner

CPGs are guidelines for best practice and are not intended as a substitute for good clinical judgment. Unusual patient presentations make it impossible to develop a CPG to match every possible clinical situation. The practitioner decides if a CPG should be applied based on patient assessment and the clinical impression. The practitioner must work in the best interest of the patient within the scope of practice for his/her clinical level on the PHECC Register. Consultation with fellow practitioners and or medical practitioners in challenging clinical situations is strongly advised.

The CPGs herein may be implemented provided:

1. The practitioner is in good standing on the PHECC practitioner's Register – Credentialed.
2. The practitioner is acting on behalf of a Licensed CPG Provider (paid or voluntary) – Licensed.
3. The practitioner is privileged by the Licensed CPG Provider on whose behalf he/she is acting to implement the specific CPG – Privileged.
4. The practitioner has received training on, and is competent in, the skills and medications specified in the CPG being utilised.

The medication dose specified on the relevant CPG shall be the definitive dose in relation to practitioner administration of medications. The principle of titrating the dose to the desired effect shall be applied. The onus rests on the practitioner to ensure that he/she is using the latest versions of CPGs, which are available on the PHECC website www.phecc.ie

Definitions

Adult	A patient of 16 years or greater, unless specified on the CPG
Child	A patient between 1 and less than or equal to (\leq) 15 years old, unless specified on the CPG
Infant	A patient between 4 weeks and less than 1 year old, unless specified on the CPG
Neonate	A patient less than 4 weeks old, unless specified on the CPG
Paediatric patient	Any child, infant or neonate

CPGs and the pre-hospital emergency care team

The aim of pre-hospital emergency care is to provide a comprehensive and coordinated approach to patient care management, thus providing each patient with the most appropriate care in the most efficient time frame.

In Ireland today, the provision of emergency care comes from a range of disciplines and includes responders (Cardiac First Responders, First Aid Responders and Emergency First Responders) and practitioners (Emergency Medical Technicians, Paramedics, Advanced Paramedics, Nurses and Doctors) from the statutory, private, auxiliary and voluntary services.

CPGs set a consistent standard of clinical practice within the field of pre-hospital emergency care. By reinforcing the role of the practitioner, in the continuum of patient care, the chain of survival and the golden hour are supported in medical and traumatic emergencies respectively.

CPGs guide the practitioner in assessment, treatment and disposition of patients who present with an acute illness or injury.

CPGs presume no intervention has been applied, nor medication administered, prior to the arrival of the practitioner. In the event of another practitioner or responder initiating care during an acute episode, the practitioner must be cognisant of interventions applied and medication doses already administered and act accordingly.

In this care continuum, the duty of care is shared among all responders/practitioners of whom each is accountable for his/her own actions. The most qualified responder/practitioner on the scene shall take the role of clinical lead. Explicit handover between responders/practitioners is essential and will eliminate confusion regarding the responsibility for care.

When a practitioner of higher clinical level on scene deems it appropriate to take clinical lead, he/she should calmly state: "My name is xx, I am an AP/P/EMT, I am assuming clinical lead."

If the practitioner of higher clinical level on scene wishes to hand over clinical lead to another practitioner (who may be of equal or lower clinical level), he/she states to the practitioner: "My name is xx, I am an AP/P/EMT, you are now clinical lead."

The practitioner acknowledges immediately and accepts clinical lead. "I am now clinical lead"

A clinical lead exchange should be recorded on the PCR in the 'continuity of care' section. There should never be any doubt as to who the clinical lead is on scene.

In the absence of a more qualified practitioner, the practitioner providing care during transport shall be designated the clinical lead as soon as practical.

Classification of CPGs

The Taxonomy for Pre-Hospital Emergency Care CPGs has changed to a new method for configuring PHECC CPGs. There are now seventeen categories developed to group common themes and categories together.

Basic Life Support – ILCOR 2020

Basic life support CPGs contained within this publication are in accordance with International Liaison Committee on Resuscitation (ILCOR) guidelines 2020.

SECTION 1 PRINCIPLES OF GENERAL CARE ... 16	8.3 External Haemorrhage – Adult 41
1.2 Primary Survey Medical – Adult 18	8.4 Harness Induced Suspension Trauma 42
1.3 Primary Survey Trauma – Adult 19	8.5 Head Injury – Adult..... 43
1.5 Secondary Survey Medical – Adult 20	8.6 Limb Injury – Adult 44
1.6 Secondary Survey Trauma – Adult 21	8.8 Spinal Injury Management 45
SECTION 2 AIRWAY AND BREATHING 22	8.9 Submersion/Immersion Incident 46
2.1 Foreign Body Airway Obstruction – Adult 22	SECTION 9 ENVIRONMENTAL 47
2.2 Advanced Airway Management – Adult 23	9.1 Hypothermia 47
2.3 Abnormal Work of Breathing – Adult 24	9.2 Heat Related Emergency – Adult 48
2.4 Exacerbation of COPD 25	SECTION 10 TOXICOLOGY 49
2.5 Asthma – Adult 26	10.1 Allergic Reaction/Anaphylaxis – Adult 49
2.7 Emergency Tracheostomy Management 27	10.2 Poisons – Adult 50
SECTION 3 CARDIAC 28	SECTION 11 INFECTIOUS 51
3.1 Acute Coronary Syndrome 28	11.1 Sepsis – Adult 51
3.2 Symptomatic Bradycardia – Adult 29	SECTION 12 MATERNAL 52
SECTION 4 CIRCULATION 30	12.1 Pregnancy Related Emergencies 52
4.2 Epistaxis..... 30	12.2 Pre-Hospital Emergency Childbirth 53
SECTION 5 MEDICAL 31	12.5 Umbilical Cord Complications 54
5.2 Decompression Illness (DCI)..... 31	12.6 Post pregnancy care 55
5.3 Glycaemic Emergency – Adult 32	12.7 New-born neonatal Care and Resuscitation. 56
5.4 Sickle Cell Crisis – Adult..... 33	12.8 Neonatal Resuscitation ≤ 6 weeks 57
SECTION 6 NEUROLOGICAL 34	SECTION 13 PAEDIATRIC 58
6.1 Altered Level of Consciousness – Adult 34	13.1 Primary Survey Medical – Paediatric 58
6.2 Pain Management – Adult..... 35	13.2 Primary Survey Trauma – Paediatric 59
6.3 Seizure/Convulsion – Adult 36	13.4 Secondary Survey – Paediatric 60
6.4 Stroke 37	13.5 Foreign Body Airway Obstruction –Paediatric 61
SECTION 7 BEHAVIOURAL AND MENTAL HEALTH EMERGENCIES 38	13.7 Inadequate Ventilations – Paediatric 62
7.1 Mental Health Emergency 38	13.8 Asthma – Paediatric 63
7.2 Behavioural Emergency..... 39	13.9 Stridor – Paediatric 64
SECTION 8 TRAUMA 40	13.11 Glycaemic Emergency – Paediatric 65
8.1 Burns – Adult 40	13.12 Sickle Cell Crisis – Paediatric 66
	13.13 Pain Management – Paediatric 67
	13.14 Seizure/Convulsion – Paediatric 68

13.15 Burns – Paediatric 69

13.16 External Haemorrhage – Paediatric 70

13.17 Shock from Blood Loss – Paediatric 71

13.18 Limb Injury – Paediatric 72

13.19 Pyrexia – Paediatric 73

13.20 Septic Shock – Paediatric 74

13.21 Allergic Reaction/Anaphylaxis – Paediatric 75

13.22 Basic Life Support – Paediatric 76

13.23 VF or pulseless VT – Paediatric 77

13.24 Asystole/PEA – Paediatric 78

13.25 Symptomatic Bradycardia – Paediatric 79

13.26 Post-Resuscitation Care - Paediatric 80

SECTION 14 RESUSCITATION 81

14.1 Basic Life Support – Adult 81

14.2 VF or pulseless VT – Adult 82

14.3 Asystole – Adult 83

14.5 Pulseless Electrical Activity – Adult 84

14.6 Post-Resuscitation Care – Adult 85

14.7 Recognition of Death – Resuscitation not
Indicated 86

14.8 Team Resuscitation 87

SECTION 15 PALLIATIVE CARE 88


15.1 End of Life – DNAR 88

SECTION 16 OPERATIONS 89

16.1 Major Emergency (Major Incident) – First
Practitioners on site 89

16.2 Major Emergency (Major Incident) –
Operational Control 90

16.3 Triage Sieve 91



Principles of general care (Practitioners)

Care principles are goals of care that apply to all patients. Scene safety, standard precautions, patient assessment, primary and secondary surveys and the recording of interventions and medications on the Patient Care Report (PCR) or the Ambulatory Care Report (ACR), are consistent principles throughout the guidelines and reflect the practice of practitioners. Care principles are the foundations for risk management and the avoidance of error.


PHECC Care Principles

1. Ensure the safety of yourself, other emergency service personnel, your patients and the public.
 - 1.1. Ensure correct PPE is utilised in all situations and is compliant with latest guidance on standard, contact, droplet and airborne PPE. Place facemasks on patients when required. Handwashing and hand hygiene should be performed before and after all patient interactions. Utilise PPE checklists for correct donning and doffing procedures.
2. A person has capacity in respect to clinical decisions affecting themselves unless the contrary is shown (Assisted Decision-Making (Capacity) Act 2015).
3. Seek consent prior to initiating interventions and/or administering medications.
4. Identify and manage life-threatening conditions.
5. Ensure adequate ventilation and oxygenation.
6. Optimise tissue perfusion.
7. Make a working diagnosis, after considering differential diagnoses.
8. Provide appropriate pain relief within the scope of practice. Pain management:
 - 8.1. should not delay the diagnosis of conditions or injuries,
 - 8.2. should be implemented for all relevant patients,
 - 8.3. should commence within ten minutes on scene,
 - 8.4. goal is to reduce pain to a tolerable level,
 - 8.5. to take cognisance of immediate and short-term pain management requirements by administering appropriate combinations of analgesia.
9. Identify and manage other conditions.
10. Place the patient in the appropriate posture according to the presenting condition.
11. Ensure maintenance of normal body temperature (unless a CPG indicates otherwise).
12. Provide reassurance at all times.
13. Monitor and record patient's vital observations.

14. Maintain responsibility for patient care until handover to an appropriate practitioner.
15. Arrange transport to an appropriate medical facility, if clinically required, and in an appropriate time frame.
16. Complete a patient care record following an interaction with a patient.
17. Identify the clinical lead on scene; this shall be the most qualified practitioner on scene. In the absence of a more qualified practitioner, the practitioner providing care during transport shall be designated clinical lead as soon as practical.
18. Ambulances, medical rooms and equipment should be decontaminated as appropriate following an interaction with a patient.


Primary Survey Medical - Adult

4/5/6.1.2
Version 5, 12/2020



Primary Survey Trauma – Adult

4/5/6.1.3
Version 5, 03/2021



The primary survey is focused on establishing the patient's clinical status and only applying interventions when they are essential to maintain life. It should be completed within one minute of arrival on scene.

Suction
OPA
NPA


EMT
Special Authorisation:
EMTs having completed the BTEC course may be privileged by a licensed CPG provider to insert an NPA on its behalf

Consider
Oxygen therapy

Maximum time on scene for life-threatening trauma: ≤ 10 minutes

Secondary Survey Medical – Adult


4.1.5
Version 3, 02/2021



Reference: Sanders, M. 2001, Paramedic Textbook 2nd Edition, Mosby
 Gleadle, J. 2003, History and Examination at a glance, Blackwell Science
 Rees, JE, 2003, Early Warning Scores, World Anaesthesia Issue 17, Article 10

Secondary Survey Trauma – Adult


4.1.6
Version 2, 02/2021



<p>Markers for multi-system trauma Systolic BP < 90 Respiratory rate < 10 or > 29 Heart rate > 120 AVPU = V, P or U on scale Mechanism of Injury</p>
--

Foreign Body Airway Obstruction – Adult

4/5.2.1
Version 3, 12/2020




After each cycle of CPR open mouth and look for object.
If visible, make one attempt to remove.

Advanced Airway Management – Adult

4.2.2
Version 5, 12/2020


EMT



EMT Special Authorisation:
EMTs may use cuffed supraglottic airways subject to maintaining competence and Medical Director authorisation


Abnormal Work of Breathing – Adult

4/5/6.2.3
Version 3, 03/2021



Exacerbation of COPD


4/5/6.2.4
Version 3, 03/2021



An exacerbation of COPD is defined as;
An event in the natural course of the disease characterised by a change in the patient's baseline dyspnoea, cough and/or sputum beyond day-to-day variability sufficient to warrant a change in management. (European Respiratory Society)


Asthma – Adult

4/5/6.2.5
Version 5, 03/2021




Emergency Tracheostomy Management

4/5/6.2.7
Version 2, 03/2021



Acute Coronary Syndrome

4.3.1
Version 4, 12/2020




Symptomatic Bradycardia – Adult

4/5/6.3.2
Version 4, 01/2021



Symptomatic includes;
Acute altered mental status
Ischemic chest discomfort
Acute heart failure
Hypotension
Signs of shock

Titrate Atropine to effect (HR > 60) and non-symptomatic




1mg Adrenaline in 100mL NaCl (0.9%)
• 1 mL = 10mcg, draw up in 1mL syringe and administer as a bolus.




Epistaxis

4/5/6.4.2
Version 5, 03/2021




Decompression Illness (DCI)

4/5/6.5.2
Version 3, 12/2020



Glycaemic Emergency – Adult

4/5/6.5.3
Version 4, 03/202




Consider Glucagon IM if not already given

Check for presence of an insulin pump; turn off or remove if present.


Sickle Cell Crisis - Adult

4/5/6.5.4
Version 2, 12/2020




Altered Level of Consciousness – Adult

4.6.1
Version 2, 12/2020




Pain Management – Adult

4/5/6.6.2
Version 6, 04/2021




Seizure/Convulsion – Adult

4.6.3
Version 3, 01/2021



Stroke

4.6.4
Version 3, 01/2021




<p>F – facial weakness Can the patient smile? Has their mouth or eye drooped? Which side?</p> <p>A – arm weakness Can the patient raise both arms and maintain for 5 seconds?</p> <p>S – speech problems Can the patient speak clearly and understand what you say?</p> <p>T – time of onset</p>
--

Behavioural and Mental Health Emergencies

Mental Health Emergency


4/5.7.1
Version 3, 03/2021



Behavioural and Mental Health Emergencies


Behavioural Emergency

4/5/6.7.2
Version 3, 03/2021




Burns – Adult

4/5/6.8.1
Version 3, 01/2021



External Haemorrhage – Adult


4/5/6.8.3
Version 5, 02/2021



EMT BTEC
Special Authorisation:
EMTs, having completed the BTEC course, may be privileged by a licensed CPG provider to insert wound closure clips on its behalf


Harness Induced Suspension Trauma

4/5/6.8.4
Version 4, 01/2021




Head Injury – Adult

4.8.5
Version 3, 12/2020




Limb Injury – Adult

4/5/6.8.6
Version 6, 03/2021



Spinal Injury Management

4.8.8
Version 4, 12/2020



- High risk factors:-** any of the following;
- dangerous mechanism of injury
 - fall from a height of greater than 1 metre or 5 steps
 - axial load to the head or base of the spine
- for example;
- diving, high-speed motor vehicle collision, rollover motor accident, ejection from a motor vehicle, accident involving motorised recreational vehicle, bicycle collision, horse riding accident, pedestrian v vehicle.
 - impaired awareness (alcohol/ drug intoxication, confused /uncooperative or ALoC)
 - age 65 years or older
 - age 2 years or younger incapable of verbal communication

- Spinal injury rule in considerations;**
- any significant distracting injuries
 - impaired awareness (alcohol/ drug intoxication, confused /uncooperative or ALoC)
 - Immediate onset of spinal/ midline back pain
 - hand or foot weakness (motor issue)
 - altered or absent sensation in the hands or feet (sensory issue)
 - priapism
 - history of past spinal problems, including previous spinal surgery or conditions that predispose to instability of the spine.
 - Unable to actively rotate their neck 45 degrees to the left and right.

PHECC Spinal Injury Management Standard

- Active spinal motion restriction; using inline techniques with or without spinal injury management devices to reduce spinal column motion.

Unlikely to have a clinically significant spinal injury


Low risk factors:- any two or more of;

- involved in a minor rear-end motor vehicle collision
- comfortable in a sitting position
- ambulatory at any time since the injury
- no midline cervical spine tenderness
- no spinal column/ midline pain

And are able to actively rotate their neck 45 degrees to the left and right


Submersion/ Immersion Incident

4/5/6.8.9
Version 3, 03/2021




Hypothermia

4.9.1
Version 4, 12/2020




Heat Related Emergency – Adult

4/5/6.9.2
Version 3, 01/2021



Allergic Reaction/Anaphylaxis – Adult

4/5/6.10.1
Version 4, 03/2021



Mild
Urticaria

Moderate
Mild symptoms +
angio oedema
simple bronchospasm




Severe/ anaphylaxis
Moderate symptoms +
A,B or C compromise
(haemodynamic and or
respiratory compromise)

Adrenaline:
Repeat at 5 minute intervals prn


Poisons - Adult

4/5.10.2
Version 5, 12/2020




Sepsis – Adult

4/5/6.11.1
Version 5, 03/2021



Pregnancy Related Emergencies

4/5/6.12.1
Version 3, 01/2021




Suspect pre-eclampsia if above symptoms present and 2 elevated BP readings 15min apart.



Follow local protocol


Pre-Hospital Emergency Childbirth

4/5/6.12.2
Version 4, 01/2021




Umbilical Cord Complications

4/5/6.12.5
Version 3, 01/2021




Post Pregnancy Care
(Including miscarriage and abortion)

4/5/6.12.6
Version 4, 01/2021




New-born Neonatal Care and Resuscitation

4/5/6.12.7
Version 5, 04/2021




Neonatal Resuscitation (≤ 6 weeks)

4/5/6.12.8
Version 1, 01/2021




Primary Survey Medical – Paediatric

4/5/6.13.1
Version 7, 01/2021




Primary Survey Trauma – Paediatric

4/5/6.13.2
Version 7, 01/2021



Secondary Survey – Paediatric

4/5/6.13.4
Version 5, 01/2021



Normal ranges (ICTS)		
Age	Pulse	Respirations
0-3 months	90 – 180	30 – 60
4-6 months	80 – 160	30 – 60
7-12 months	80 – 140	25 – 45
1-3 years	75 – 130	20 – 30
4-6 years	70 – 110	16 – 24
≥ 7 years	60 – 90	14 – 20

Estimated weight
Neonate = 3.5kg
Six months = 6kg
One to five years = (age x 2) + 8kg
Greater than 5 years = (age x 3) + 7kg

Use age appropriate language for patient


Children and adolescents should always be examined with a chaperone (usually a parent) where possible

If child protection concerns are present

Report findings as per Children First Act 2015 to ED staff and Tusla in a confidential manner


Foreign Body Airway Obstruction – Paediatric

4/5.13.5
Version 4, 03/2021




Abnormal Work of Breathing - Paediatric

4/5/6.13.7
Version 4, 03/2021




Asthma – Paediatric

4/5/6.13.8
Version 4, 01/2021



Stridor – Paediatric

4/5/6.13.9
Version 5, 01/2021



Signs of Croup may include:

- Hypoxia/ cyanosis
- Marked respiratory distress
- Stridor
- Irritability or lethargy
- Marked increased respiratory rate


If persistent treat as severe croup
If symptoms are intermittent treat as moderate croup

Nebulised Adrenaline may be repeated after 30 minutes prn

Maximum Dexamethasone administered within the past 72 hours not to exceed 600mcg/kg


Glycaemic Emergency – Paediatric

4/5/6.13.11
Version 6, 01/2021




Sickle Cell Crisis – Paediatric

4/5/6.13.12
Version 2, 01/2021




Pain Management – Paediatric

4/5/6.13.13
Version 9, 03/2021




Seizure/Convulsion – Paediatric

4.13.14
Version 4, 12/2020




Burns – Paediatric

4/5/6.13.15
Version 4, 01/2021




External Haemorrhage – Paediatric

4/5/6.13.16
Version 5, 01/2021



Shock from Blood Loss – Paediatric


4.13.17
Version 3, 12/2020



Signs of inadequate perfusion
A: (not directly affected)
B: Increased respiratory rate (without increased effort)
C: Tachycardia Diminished/absent peripheral pulses Delayed capillary refill
D: Irritability/ confusion / ALoC
E: Cool extremities, mottling

Limb Injury – Paediatric


4/5/6.13.18
Version 1, 04/2021



For a limb threatening injury treat as an emergency and pre-alert ED


Pyrexia – Paediatric

4/5/6.13.19
Version 3, 01/2021




Sepsis – Paediatric

4/5/6.13.20
Version 5, 03/2021




Allergic Reaction/Anaphylaxis – Paediatric

4/5/6.13.21
Version 5, 04/2021




Basic Life Support – Paediatric

4/5/6.13.22
Version 5, 02/2021



VF or pVT – Paediatric

4/5/6.13.23
Version 6, 01/2021



Consider transport to ED if no change after 20 minutes resuscitation
If no ALS available

Defibrillation:
< 8 years use paediatric defibrillation system (if not available use adult pads)

If refractory VF/pVT post Adrenaline and 3rd shock
Amiodarone 5 mg/kg IV/IO


Special Authorisation:
Advanced Paramedics are authorised to substitute Amiodarone with a one off bolus of Lidocaine (1-1.5mg/kg IV) if Amiodarone is not available

- Consider causes and treat as appropriate:**
- Hydrogen ion acidosis
 - Hyper/ hypokalaemia
 - Hypothermia
 - Hypovolaemia
 - Hypoxia
 - Thrombosis – pulmonary
 - Tension pneumothorax
 - Thrombus – coronary
 - Tamponade – cardiac
 - Toxins
 - Trauma

* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm

Asystole/PEA – Paediatric

4/5/6.13.24
Version 5, 01/2021




- Consider causes and treat as appropriate:
- Hydrogen ion acidosis
 - Hyper/ hypokalaemia
 - Hypothermia
 - Hypovolaemia
 - Hypoxia
 - Thrombosis – pulmonary
 - Tension pneumothorax
 - Thrombus – coronary
 - Tamponade – cardiac
 - Toxins
 - Trauma

Consider fluid challenge
NaCl (0.9%) 20 mL/kg IV/IO

* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Symptomatic Bradycardia – Paediatric

4/5/6.13.25
Version 5, 01/2021




Post-Resuscitation Care – Paediatric

4.13.26
Version 4, 04/2021




Basic Life Support – Adult

4/5/6.14.1
Version 4, 02/2021



VF or pVT – Adult


4/5/6.14.2
Version 5, 01/2021



* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Asystole – Adult

4.14.3
Version 4, 12/2020



Pulseless Electrical Activity – Adult

4/5/6.14.5
Version 4, 01/2021



- Consider causes and treat as appropriate:**
- Hydrogen ion acidosis
 - Hyper/ hypokalaemia
 - Hypothermia
 - Hypovolaemia
 - Hypoxia
 - Thrombosis – pulmonary
 - Tension pneumothorax
 - Thrombus – coronary
 - Tamponade – cardiac
 - Toxins
 - Trauma


Consider fluid challenge
NaCl (0.9%) 1L IV/IO
Repeat prn

If Tricyclic antidepressant toxicity or harness induced suspension trauma consider
Sodium Bicarbonate (8.4%) 1 mEq/kg IV/IO

* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Post-Resuscitation Care – Adult

4.14.6
Version 4, 03/2021




Recognition of Death – Resuscitation not Indicated

4.14.7
Version 2, 12/2020




Team Resuscitation

4/5/6.14.8
Version 2, 03/2021



End of Life – DNAR

4.15.1
Version 2, 12/2020




Major Emergency (Major Incident) – First Practitioners on site

4/5/6.16.1
Version 3, 12/2020



Irish (Major Emergency) terminology in black
UK (Major Incident) terminology in blue



If single Practitioner is first on site combine both roles until additional Practitioners arrive

METHANE message
 M – Major Emergency declaration / standby
 E – Exact location of the emergency
 T – Type of incident (transport, chemical etc.)
 H – Hazards present and potential
 A – Access / egress routes
 N – Number of casualties (injured or dead)
 E – Emergency services present and required

The first ambulance crew does not provide care or transport of patients as this interferes with their ability to liaise with other services, to assess the scene and to provide continuous information as the incident develops


Major Emergency (Major Incident) – Operational Control

4/5/6.16.2
Version 3, 12/2020



Irish (Major Emergency) terminology in black
UK (Major Incident) terminology in blue

If Danger Area identified, entry to Danger Area is controlled by a Senior Fire Officer or an Garda Síochána



Entry to Inner Cordon (Bronze Area) is limited to personnel providing emergency care and/or rescue
Personal Protective Equipment required

Entry to Outer Cordon (Silver area) is controlled by an Garda Síochána

Management structure for; Outer Cordon, Tactical Area (Silver Area)
On-Site Co-ordinator
HSE Controller of Operations (Ambulance Incident Officer)
Site Medical Officer (Medical Incident Officer)
Local Authority Controller of Operations (Fire Incident Officer)
Garda Controller of Operations (Police Incident Officer)

Management structure for; Inner Cordon, Operational Area (Bronze Area)
Forward Ambulance Incident Officer (Forward Ambulance Incident Officer)
Forward Medical Incident Officer (Forward Medical Incident Officer)
Fire Service Incident Commander (Forward Fire Incident Officer)
Garda Cordon Control Officer (Forward Police Incident Officer)


Please note that Controller of Operations may be other than ambulance or fire officers, depending on the nature of the emergency

Other management functions for; Major Emergency site
Casualty Clearing Officer
Triage Officer
Ambulance Parking Point Officer
Ambulance Loading Point Officer
Communications Officer
Safety Officer



Triage Sieve

4/5/6.16.3
Version 2, 12/2020



Triage is a dynamic process

Medication Formulary for Emergency Medical Technician

The Medication Formulary is published by the Pre-Hospital Emergency Care Council (PHECC) to enable pre-hospital emergency care practitioners to be competent in the use of medications permitted under Medicinal Products 7th Schedule (SI 300 of 2014).

This is a summary document only and practitioners are advised to consult with official publications to obtain detailed information about the medications used.

The Medication Formulary is recommended by the Medical Advisory Committee (MAC) prior to publication by Council.

The medications herein may be administered provided:

1. The practitioner is in good standing on the PHECC practitioner's Register.
2. The practitioner complies with the Clinical Practice Guidelines (CPGs) published by PHECC.
3. The practitioner is acting on behalf of an organisation (paid or voluntary) that is a PHECC licensed CPG provider.
4. The practitioner is privileged, by the organisation on whose behalf he/she is acting, to administer the medications.
5. The practitioner has received training on, and is competent in, the administration of the medication.
6. The medications are listed on the Medicinal Products 7th Schedule.

The context for administration of the medications listed here is outlined in the CPGs.

Every effort has been made to ensure accuracy of the medication doses herein. The dose specified on the relevant CPG shall be the definitive dose in relation to practitioner administration of medications. The principle of titrating the dose to the desired effect shall be applied. The onus rests on the practitioner to ensure that he/she is using the latest versions of CPGs which are available on the PHECC website www.phecc.ie

Sodium Chloride 0.9% (NaCl) is the IV/IO fluid of choice for pre-hospital emergency care.

Water for injection shall be used when diluting medications, however if not available NaCl (0.9%) may be used if not contraindicated.

All medication doses for patients ≤ 15 years shall be calculated on a weight basis unless an age-related dose is specified for that medication.

The route of administration should be appropriate to the patient's clinical presentation. IO access is authorised for advanced paramedics for life threatening emergencies (or under medical direction).

The dose for paediatric patients may never exceed the adult dose.

Approved Paediatric weight estimations are:

Neonate =	3.5 Kg
Six months =	6 Kg
One to five years =	(age x 2) + 8 Kg
Greater than 5 years =	(age x 3) + 7 Kg

Pregnancy caution:

Medications should be prescribed in pregnancy only if the expected benefit to the mother is thought to be greater than the risk to the foetus, and all medications should be avoided, if possible, during the first trimester.

PHECC practitioners therefore should avoid using medications in early pregnancy unless absolutely essential and where possible medical advice should be sought prior to administration.

Paramedic authorisation for IV infusion continuation

PHECC registered paramedics are authorised to continue an established IV infusion in the absence of an advanced paramedic or doctor during transportation.

Medication Formulary Age Designations

Index of medication formulary (Adult \geq 16 and Paediatric \leq 15 unless otherwise stated)

This version contains 14 medications.

Please visit www.phecc.ie for the latest edition/version

Amendments to the Emergency Medical Technician 2017 Edition:

New Medications introduced:

- Activated Charcoal

Changes to Monographs

1. Class and Description headings have merged to one Classification heading in line with BNF drug descriptors
2. Long term side effects have been removed unless essential
3. Pharmacology/Action has been removed unless essential information

EPINEPHRINE (1:1,000) CHANGES TO ADRENALINE (1:1000)

Heading	Add	Delete								
Medication	Adrenaline 1:1000.	Epinephrine 1:1000.								
Indications	Stridor, Symptomatic Bradycardia and Cardiogenic Shock.									
Contra-indications	Hypersensitivity to excipients.									
Usual Dosages	<table border="0"> <tr> <td>< 6 months</td> <td>10 mcg/kg IM</td> </tr> <tr> <td>6 months to < 6 years</td> <td>150 mcg (0.15 mL IM)</td> </tr> <tr> <td>≥ 6 years to < 12 years</td> <td>300 mcg (0.3 mL IM)</td> </tr> <tr> <td>≥ 12 years</td> <td>300 mcg (0.3 mL) (if child small or prepubital) or 500 (0.5 mL IM)</td> </tr> </table>	< 6 months	10 mcg/kg IM	6 months to < 6 years	150 mcg (0.15 mL IM)	≥ 6 years to < 12 years	300 mcg (0.3 mL IM)	≥ 12 years	300 mcg (0.3 mL) (if child small or prepubital) or 500 (0.5 mL IM)	<p>All dosing which was previously recommended under the following age categories</p> <p>< 6 months, 6 months to 5 years, 6 to 8 years, > 8 years.</p>
< 6 months	10 mcg/kg IM									
6 months to < 6 years	150 mcg (0.15 mL IM)									
≥ 6 years to < 12 years	300 mcg (0.3 mL IM)									
≥ 12 years	300 mcg (0.3 mL) (if child small or prepubital) or 500 (0.5 mL IM)									

ASPIRIN		
Heading	Add	Delete
Classification	Merge Class and Description to Classification: Antithrombotic – Antiplatelet Drug which reduces clot formation.	Class. Description.
Description		Anti-inflammatory agent and an inhibitor of platelet function. Useful agent in the treatment of various thromboembolic diseases such as acute myocardial infarction.
Pharmacology/ Action		Antithrombotic: Inhibits the formation of thromboxane A2, which stimulates platelet aggregation and artery constriction. This reduces clot/thrombus formation in an MI.
Long term side-effects		Generally mild and infrequent but incidence of gastro-intestinal irritation with slight asymptomatic blood loss, increased bleeding time, bronchospasm and skin reaction in hypersensitive patients.

CHLORPHENAMINE																												
Heading	Add	Delete																										
Classification	Sedating antihistamine – H2 receptor antagonists.	Class: Antihistamine. Description: H1 antagonist to counteract the effects of histamine release.																										
Usual dosages	<p>For IV route, administer over 1 minute. May dilute with Sodium Chloride 0.9% for convenient administration volume of small doses.</p> <table border="1"> <thead> <tr> <th>Severity</th> <th>Age</th> <th>Dose and route of administration</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Mild</td> <td>6 to 11 years</td> <td>2 mg PO (EMT / P / AP)</td> </tr> <tr> <td>≥ 12 years</td> <td>4 mg PO (EMT / P / AP)</td> </tr> <tr> <td rowspan="4">Moderate</td> <td>1 month – 6 months</td> <td>0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)</td> </tr> <tr> <td>>6 months - < 6 years</td> <td>2.5 mg IM (EMT / P) or 2.5 mg IV (AP)</td> </tr> <tr> <td>6 to < 12 years</td> <td>2 mg PO or 5 mg IM (EMT / P) or 5 mg IV (AP).</td> </tr> <tr> <td>≥ 12 years</td> <td>4 mg PO or 10 mg IM (EMT / P) or 10 mg IV (AP)</td> </tr> <tr> <td rowspan="4">Severe</td> <td>1 month - <6 months</td> <td>0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)</td> </tr> <tr> <td>>6 months - <6 years</td> <td>2.5 mg IM (EMT / P) or 2.5 mg IV (AP)</td> </tr> <tr> <td>6 to < 12 years</td> <td>5 mg IM (EMT / P) or 5 mg IV (AP)</td> </tr> <tr> <td>≥ 12 years</td> <td>10 mg IM (EMT / P) or 10 mg IV (AP)</td> </tr> </tbody> </table>	Severity	Age	Dose and route of administration	Mild	6 to 11 years	2 mg PO (EMT / P / AP)	≥ 12 years	4 mg PO (EMT / P / AP)	Moderate	1 month – 6 months	0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)	>6 months - < 6 years	2.5 mg IM (EMT / P) or 2.5 mg IV (AP)	6 to < 12 years	2 mg PO or 5 mg IM (EMT / P) or 5 mg IV (AP).	≥ 12 years	4 mg PO or 10 mg IM (EMT / P) or 10 mg IV (AP)	Severe	1 month - <6 months	0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)	>6 months - <6 years	2.5 mg IM (EMT / P) or 2.5 mg IV (AP)	6 to < 12 years	5 mg IM (EMT / P) or 5 mg IV (AP)	≥ 12 years	10 mg IM (EMT / P) or 10 mg IV (AP)	Removal of all existing paediatric dosing.
Severity	Age	Dose and route of administration																										
Mild	6 to 11 years	2 mg PO (EMT / P / AP)																										
	≥ 12 years	4 mg PO (EMT / P / AP)																										
Moderate	1 month – 6 months	0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)																										
	>6 months - < 6 years	2.5 mg IM (EMT / P) or 2.5 mg IV (AP)																										
	6 to < 12 years	2 mg PO or 5 mg IM (EMT / P) or 5 mg IV (AP).																										
	≥ 12 years	4 mg PO or 10 mg IM (EMT / P) or 10 mg IV (AP)																										
Severe	1 month - <6 months	0.25 mg/kg IM (EMT / P) or 0.25 mg/kg IV (AP)																										
	>6 months - <6 years	2.5 mg IM (EMT / P) or 2.5 mg IV (AP)																										
	6 to < 12 years	5 mg IM (EMT / P) or 5 mg IV (AP)																										
	≥ 12 years	10 mg IM (EMT / P) or 10 mg IV (AP)																										
Additional information		For IV route, administer over 1 minute. May dilute with Sodium Chloride 0.9% for convenient administration volume of small doses.																										
Side-effects	Reworded: Causes drowsiness, do not drive or operate machinery.																											

GLUCAGON

Heading	Add	Delete
Usual dosages	Paediatric: ≥ 1 month and < 25 kg: 500 mcg IM. ≥ 1 month and ≥ 25 kg: 1 mg IM.	Paediatric: 1 - 8 years - 0.5 mg (500 mcg) IM. 8 years - 1 mg IM.
Side-effects	Common: Nausea Uncommon: Vomiting. Rare: may cause hypotension/ dizziness/ headache.	

GLUCOSE GEL

Heading	Add	Delete
Classification	Class and Description merged.	Class. Description.
Administration	CPG 4/5/6.12.7: New-born Neonatal Care and Resuscitation.	

GLYCERYL TRINITRATE (GTN)

Heading	Add	Delete
Classification		Class. Description.
Presentation		(0.4 mg).
Usual Dosages	Angina or MI: 400 mcg sublingual. (Repeat at 3-5 min intervals, Max: 1200 mcg). EFR: assist administration - 400 mcg sublingual max. Pulmonary oedema: 800 mcg / 2 sprays (repeat x 1 PRN) (P & AP).	0.4 mg. 1.2 mg. 0.4 mg. 0.8 mg.
Pharmacology / Action		Remove complete section.

IBUPROFEN

Heading	Add	Delete
Classification	Analgesics: Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Pain and Inflammation in musculoskeletal disorders.	Class: Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Description: It is an anti-inflammatory analgesic.
Contra-Indications	Body weight < 5 kg.	
Long term side-effects		Remove list of long-term side-effects.

METHOXYFLURANE

Heading	Add	Delete
Classification	Anaesthetics. General: Volatile anaesthetic agent.	
Contra-Indications	Malignant Hyperthermia.	

NALOXONE

Heading	Add	Delete
Usual Dosages	400 mcg. 800 mcg.	0.4 mg. 0.8 mg.

NITROUS OXIDE 50% AND OXYGEN 50%

Heading	Add	Delete
Additional Information	Caution should be issued before using Entonox with patients who have known Chronic Obstructive Pulmonary Disease (COPD) or other conditions where compromised chemoreceptor sensitivity/function may be present. May cause respiratory depression and increases in PaCO ₂ . In cold temperatures warm cylinder and invert at least 3 times to ensure mix of gases. Prolonged or frequent use of ENTONOX may result in megaloblastic marrow changes, myeloneuropathy and sub-acute combined degeneration of the spinal cord.	In cold temperatures warm cylinder and invert to ensure mix of gases.

OXYGEN		
Heading	Add	Delete
Clinical Level		
Classification	Merged Class and Description.	Class. Description.
Pharmacology/Action		Pharmacology/Action Oxygenation of tissue/organs.
Additional Information	Caution with emollients containing paraffin e.g. lip balms & moisturisers – may lead to skin burns.	

PARACETAMOL		
Heading	Add	Delete
Presentation	500 mg of paracetamol in 50 mL solution for infusion.	0.1 mg.
Usual Dosages	15 mg/kg PO. PR (AP). > 1 month < 1 year - 80 mg PR.	20 mg/kg PO. > 1 month < 1 year - 90 mg PR.
Side effects		Long term side-effects.

SALBUTAMOL		
Heading	Add	Delete
Classification	Beta-2 Adrenoceptor agonist selective – short acting.	Class: Sympathetic agonist. Description: Sympathomimetic that is selective for Beta-2 Adrenergic receptors.
Presentation	100 mcg.	0.1 mg.
Usual Dosages	100 mcg metered aerosol spray.	0.1 mg metered aerosol spray.
Pharmacology / Action		Remove text/section Beta-2 agonist/ Bronchodilation/ relaxation of smooth muscle.

Clinical Level:



MEDICATION	ACTIVATED CHARCOAL
Classification	Antidotes and Chelators – Intestinal adsorbents: reduction of absorption of poisons in the GI system / active elimination of poisons.
Presentation	Activated charcoal granules for suspension.
Administration	Oral suspension (PO). (CPG: 6.10.2).
Indications	Emergency treatment of acute oral poisoning or drug overdose.
Contra-Indications	Although activated charcoal is not contraindicated in poisoning by strong acids and alkalis and other corrosive substances, its value as a detoxicant for these substances is limited. Activated Charcoal is poor in binding cyanide, iron salts and some solvents including methanol, ethanol and ethylene glycol.
Usual Dosages	Adult: 50g PO. Reconstitute with water as directed by manufacturer. The reconstituted product should be taken immediately. Repeat as necessary. Paediatric: Not Indicated.
Side effects	Bezoar/ Constipation/ diarrhoea/ GI disorders/ Black stools. Caution: aspiration may lead to airway obstruction.
Additional information	May be mixed with soft drinks or fruit juice for ease of administration & to mask the taste. Substances which may be absorbed by Activated charcoal (but are not limited to) include: Aspirin & salicylates/ Barbiturates/ Benzodiazepines/ Chlormethiazole/ Chloroquine/ Chlorpromazine & related phenothiazines/ Clonidine/ Cocaine and other stimulants/ Digoxin and digitoxin/ Ibuprofen/ Mefenamic acid/ Mianserin/ Nicotine/ Paracetamol/ Paraquat/ Phenelzine and other MAOIs/ Phenytoin/ Propranolol and other Beta Blockers/ Quinine/ Theophylline/ Zidovudine.

Clinical Level:



MEDICATION		ADRENALINE (1:1,000)								
Classification	Sympathetic agonist, Sympathomimetic – Vasoconstrictor. Acts on both alpha & beta receptors and increases both heart rate and contractility. It can cause peripheral vasodilation (beta) or vasoconstriction (alpha).									
Presentation	Pre-filled syringe, ampoule or Auto injector. 1 mg/1 mL (1:1,000).									
Administration	Intramuscular (IM), Intravenous (IV) and Nebulisation (Neb). (CPG: 2/3.10.1 2/3.13.21, 4/5/6.3.2, 4/5/6.10.1, 4/5/6.11.1, 4/5/6.13.9, 5/6.13.20, 4/5/6.13.21, 5/6.14.6)									
Indications	Severe allergic reaction/ anaphylaxis, Stridor, Symptomatic Bradycardia and Cardiogenic shock.									
Contra-Indications	Hypersensitivity to excipients.									
Usual Dosages	<p>Adult: Anaphylaxis 500mcg IM (0.5mL of 1: 1,000).</p> <p>EFR assist patient – 0.3 mg (Auto injector). (Repeat every 5 minutes PRN).</p> <p>Adult: Symptomatic Bradycardia / Cardiogenic shock: 10mcg IV/IO repeat PRN. (Dilute 1 mg Adrenaline in 100 mL NaCl and draw up in 1 mL syringe, administer the dose over 1 minute). (Off-license).</p> <p>Anaphylaxis Paediatric:</p> <table border="1"> <tr> <td><6 months</td> <td>10 mcg/kg IM</td> </tr> <tr> <td>6 months to < 6 years</td> <td>150 mcg (0.15 mL IM)</td> </tr> <tr> <td>≥ 6 years to < 12 years</td> <td>300 mcg (0.3 mL IM)</td> </tr> <tr> <td>≥ 12 years</td> <td>300 mcg (0.3 mL) (if child small or prepubital) or 500 mcG (0.5 mL IM)</td> </tr> </table> <p>EFR assist patient –</p> <p>6 months < 10 years: 0.15 mg (Auto injector) (repeat every 5 minutes PRN). ≥ 10 years: 0.3 mg (Auto injector) (repeat every 5 minutes PRN).</p> <p>Stridor (P/ AP):</p> <p>< 1 Year: 2.5 mg NEB. ≥ 1 year: 5 mg NEB (repeat after 30 minutes PRN) (AP).</p> <p>Sepsis (AP): Adrenaline 0.1 mcg/kg IV/IO.</p>		<6 months	10 mcg/kg IM	6 months to < 6 years	150 mcg (0.15 mL IM)	≥ 6 years to < 12 years	300 mcg (0.3 mL IM)	≥ 12 years	300 mcg (0.3 mL) (if child small or prepubital) or 500 mcG (0.5 mL IM)
<6 months	10 mcg/kg IM									
6 months to < 6 years	150 mcg (0.15 mL IM)									
≥ 6 years to < 12 years	300 mcg (0.3 mL IM)									
≥ 12 years	300 mcg (0.3 mL) (if child small or prepubital) or 500 mcG (0.5 mL IM)									
Side effects	Palpitations / Tachyarrhythmias / Hypertension / Angina-like symptoms.									
Additional information	N.B. Double check the concentration on pack before use.									

Clinical Level:



MEDICATION	ASPIRIN
Classification	Antithrombotic – Antiplatelet Drug which reduces clot formation.
Presentation	300 mg dispersible tablet. 300 mg Enteric Coated (EC) tablet.
Administration	Orally (PO) - dispersed in water, or to be chewed if not dispersible form. (CPG: 5/6.3.1, 4.3.1, 1/2/3.3.1).
Indications	Cardiac chest pain or suspected myocardial infarction. Management of unstable angina and non ST-segment elevation myocardial infarction (NSTEMI). Management of ST-segment elevation myocardial infarction (STEMI).
Contra-Indications	Active symptomatic gastrointestinal (GI) ulcer/ Bleeding disorder (e.g. haemophilia)/ Known severe adverse reaction/ Patients < 16 years old (risk of Reye's Syndrome).
Usual Dosages	Adult: 300 mg Tablet. Paediatric: Contraindicated.
Side effects	Epigastric pain and discomfort/ Bronchospasm/ Gastrointestinal haemorrhage/ Increased bleeding times/ skin reactions in hypersensitive patients.
Additional information	Aspirin 300 mg is indicated for cardiac chest pain, regardless if patient is on an anti-coagulant or is already on Aspirin. If the patient has swallowed Aspirin EC (enteric coated) preparation without chewing, the patient should be regarded as not having taken any Aspirin; administer 300 mg PO.

Clinical Level:



MEDICATION	CHLORPHENAMINE																										
Classification	Sedating Antihistamine – H2 receptor antagonist.																										
Presentation	10 mg in 1 mL ampoule. 4 mg tablet.																										
Administration	Intravenous (IV), Intramuscular (IM) and Orally (PO). (CPG: 4/5/6.10.1, 4/5/6.13.21).																										
Indications	Anaphylaxis or allergic reaction.																										
Contra-Indications	Known severe adverse reaction/ Pre-coma states.																										
Usual Dosages	<p style="text-align: center;">For IV route, administer over 1 minute</p> <p style="text-align: center;">IV: May dilute with Sodium Chloride 0.9% for convenient administration volume of small doses.</p> <p><i>Adult:</i></p> <p><i>Allergic reaction</i></p> <p><i>Mild: 4 mg PO (EMT / P / AP).</i></p> <p><i>Moderate: 4 mg PO or 10 mg IM (EMT / P) or 10 mg IV (AP).</i></p> <p><i>Severe/Anaphylaxis: 10 mg IM (EMT / P) or 10 mg IV (AP).</i></p> <p><i>Paediatric:</i></p> <table border="1"> <thead> <tr> <th>Severity</th> <th>Age</th> <th>Dose and route of administration</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Mild</td> <td>6 to 11 years</td> <td>2 mg PO (EMT/ P/ AP)</td> </tr> <tr> <td>≥ 12 years</td> <td>4 mg PO (EMT/ P /AP)</td> </tr> <tr> <td rowspan="4">Moderate</td> <td>1 month - 6 months</td> <td>0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)</td> </tr> <tr> <td>> 6 months - < 6 years</td> <td>2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)</td> </tr> <tr> <td>6 to < 12 years</td> <td>2 mg PO or 5 mg IM (EMT/ P) or 5 mg IV (AP).</td> </tr> <tr> <td>≥ 12 years</td> <td>4 mg PO or 10 mg IM (EMT/ P) or 10 mg IV (AP)</td> </tr> <tr> <td rowspan="4">Severe</td> <td>1 month - 6 months</td> <td>0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)</td> </tr> <tr> <td>> 6 months - < 6 years</td> <td>2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)</td> </tr> <tr> <td>6 to < 12 years</td> <td>5 mg IM (EMT/ P) or 5 mg IV (AP)</td> </tr> <tr> <td>≥ 12 years</td> <td>10 mg IM (EMT/ P) or 10 mg IV (AP)</td> </tr> </tbody> </table>	Severity	Age	Dose and route of administration	Mild	6 to 11 years	2 mg PO (EMT/ P/ AP)	≥ 12 years	4 mg PO (EMT/ P /AP)	Moderate	1 month - 6 months	0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)	> 6 months - < 6 years	2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)	6 to < 12 years	2 mg PO or 5 mg IM (EMT/ P) or 5 mg IV (AP).	≥ 12 years	4 mg PO or 10 mg IM (EMT/ P) or 10 mg IV (AP)	Severe	1 month - 6 months	0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)	> 6 months - < 6 years	2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)	6 to < 12 years	5 mg IM (EMT/ P) or 5 mg IV (AP)	≥ 12 years	10 mg IM (EMT/ P) or 10 mg IV (AP)
Severity	Age	Dose and route of administration																									
Mild	6 to 11 years	2 mg PO (EMT/ P/ AP)																									
	≥ 12 years	4 mg PO (EMT/ P /AP)																									
Moderate	1 month - 6 months	0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)																									
	> 6 months - < 6 years	2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)																									
	6 to < 12 years	2 mg PO or 5 mg IM (EMT/ P) or 5 mg IV (AP).																									
	≥ 12 years	4 mg PO or 10 mg IM (EMT/ P) or 10 mg IV (AP)																									
Severe	1 month - 6 months	0.25 mg/kg IM (EMT/ P) or 0.25 mg/kg IV (AP)																									
	> 6 months - < 6 years	2.5 mg IM (EMT/ P) or 2.5 mg IV (AP)																									
	6 to < 12 years	5 mg IM (EMT/ P) or 5 mg IV (AP)																									
	≥ 12 years	10 mg IM (EMT/ P) or 10 mg IV (AP)																									
Side effects	Causes drowsiness, do not drive or operate machinery.																										
Additional information	Use with caution in epilepsy/ Prostatic hypertrophy/ Glaucoma/ Hepatic disease/ Bronchitis/ Bronchiectasis/ Thyrotoxicosis/ Raised intra-ocular pressure/ Severe hypertension/ Cardiovascular disease/ Bronchial asthma.																										

Clinical Level:



MEDICATION	GLUCAGON
Classification	Hypoglycaemia: Glycogenolytic Hormones.
Presentation	1 mg vial powder and solution for reconstitution (1 mL).
Administration	Intramuscular (IM). (CPG: 4/5/6.5.3, 4/5/6.13.11).
Indications	Hypoglycaemia in patients unable to take oral glucose or unable to gain IV access, with a blood glucose level < 4 mmol/L.
Contra-Indications	< 1 month/ Phaeochromocytoma/ Known Severe Adverse Reactions
Usual Dosages	Adult: 1 mg IM. Paediatric: ≥ 1 month and < 25kg: 500 mcg IM. ≥ 1 month and ≥ 25kg: 1 mg IM.
Side effects	Common: Nausea. Uncommon: Vomiting. Rare: may cause Hypotension/ Dizziness/ Headache.
Additional information	May be ineffective in patients with low stored glycogen e.g. prior use in previous 24 hours, alcohol dependent patients with liver disease. Store in refrigerator. Stable at room temperature for 18 months, use immediately once reconstituted. Protect from light. Hypoglycaemic paediatric patients who are not diagnosed as diabetic should not be administered Glucagon. (this does not preclude the administration of glucose gel or glucose solution to treat hypoglycaemia).

Clinical Level:



MEDICATION	GLUCOSE GEL
Classification	Nutrients. Sugars: Antihypoglycaemic.
Presentation	Glucose gel in a tube or sachet.
Administration	Buccal administration: Administer gel to the inside of the patient's cheek and gently massage the outside of the cheek. (CPG: 4/5/6.5.3, 4/5/6.12.7 4/5/6.13.11).
Indications	Hypoglycaemia. Blood glucose < 4 mmol/L.
Contra-Indications	Known severe adverse reaction.
Usual Dosages	Adult: 10 – 20 g buccal (Recheck blood glucose and repeat after 15 min if required). Paediatric: New-born neonate 2 - 4 mL if blood glucose ≤ 2.6 mmol/L. ≤ 8 years 5 – 10 g buccal (recheck blood glucose and repeat after 15 mins if required). > 8 years 10 – 20 g buccal (recheck blood glucose and repeat after 15 mins if required).
Side effects	May cause vomiting in patients under the age of 5 years if administered too quickly.
Additional information	Glucose gel will maintain glucose levels once raised but should be used secondary to Dextrose to reverse hypoglycaemia. Proceed with caution: Patients with airway compromise. Altered level of consciousness.

Clinical Level:



MEDICATION	GLYCERYL TRINITRATE (GTN)
Classification	Nitrate. Potent coronary vasodilator/ reduces BP/ Dilation of systemic veins.
Presentation	<i>Aerosol spray</i> : Metered dose of 400 mcg.
Administration	<i>Sublingual</i> : Hold the pump spray vertically with the valve head uppermost. Place as close to the mouth as possible and spray under the tongue. The mouth should be closed immediately after each dose. (CPG: 4/5/6.2.6, 4/5/6.3.1, 1/2/3.3.1).
Indications	Angina/ suspected myocardial infarction (MI). <i>EFR</i> : may assist with administration. <i>EMT</i> : Angina/ suspected myocardial infarction (MI) with systolic BP \geq 110 mmHg. <i>Advanced Paramedics and Paramedics</i> - Pulmonary oedema
Contra-Indications	SBP < 90 mmHg/ Viagra or other phosphodiesterase type 5 inhibitors (Sildenafil, Tadalafil and Vardenafil) used within previous 24 hours/ Severe mitral stenosis/ Known severe adverse reaction.
Usual Dosages	<i>Adult</i> : <i>Angina or MI</i> : 400 mcg sublingual. (Repeat at 3-5 min intervals, Max: 1200 mcg). <i>EFR</i> : assist administration - 400 mcg sublingual max. <i>Pulmonary oedema</i> : 800 mcg/ 2 sprays (repeat x 1 PRN) (P & AP). <i>Paediatric</i> : Not indicated.
Side effects	Headache/ Transient Hypotension/ Flushing/ Dizziness.
Additional information	Caution with inferior wall MI with right ventricular involvement as this may lead to profound hypotension. If the pump is new or it has not been used for a week or more the first spray should be released into the air.

Clinical Level:



MEDICATION	IBUPROFEN
Classification	Analgesics: Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Pain and Inflammation in musculoskeletal disorders.
Presentation	Suspension 100 mg in 5 mL and 200 mg in 5 mL. 200 mg, 400 mg tablets.
Administration	Orally (PO). (CPG: 4/5/6.6.2, 4/5/6.13.13).
Indications	Mild to moderate pain.
Contra-Indications	Not suitable for children under 3 months (or body weight <5kg)/ Patient with history of asthma exacerbated by Aspirin/ Pregnancy/ Peptic ulcer disease/ Known renal failure/ Known severe liver failure/ Known severe heart failure/ Concurrent NSAID use (e.g. Diclofenac, Naproxen)/ Known severe adverse reaction.
Usual Dosages	Adult: 400 mg PO (Mild pain). 600 mg PO (Moderate pain). Paediatric: 10 mg/kg PO to a maximum of 400 mg.
Side effects	Skin rashes/ Gastrointestinal intolerance and bleeding.
Additional information	If Ibuprofen administered in previous 6 hours, adjust the dose downward by the amount given by other sources resulting in a maximum of 10 mg/Kg or 400 mg for paediatrics. Caution with significant burns or poor perfusion due to risk of kidney failure. Caution if on oral anticoagulant (e.g. Warfarin, Rivaroxaban, Apixaban, Edoxaban) due to increased bleeding risk. Ibuprofen may be combined with Paracetamol for synergic effect.

Clinical Level:



MEDICATION	METHOXYFLURANE
Classification	Anaesthetics. General: Volatile anaesthetic agent.
Presentation	3 mL vial with a tear off tamper-evident seal which is administered via carbon inhalation vapouriser.
Administration	Inhaled (INH) through an activated Carbon Chamber (self-administered). (CPG: 4/5/6.6.2, 4/5/6.13.13).
Indications	Adult: Moderate to severe pain. Paediatric: Moderate to severe pain.
Contra-Indications	< 5 years old Altered LOC due to head injury, drugs or alcohol/ Cardiovascular instability/ Respiratory depression/ Renal Failure or Impairment/ Known Severe Adverse Reactions/ Malignant Hyperthermia.
Usual Dosages	Adult: 3 mL (INH) (repeat x 1 only PRN). Paediatric: 3 mL (INH) (repeat x 1 only PRN).
Side effects	Amnesia/ Anxiety/ Depression/ Dizziness/ Dysarthria/ Dysgeusia/ Euphoria/ Headache/ Sensory neuropathy/ Somnolence/ Hypotension/ Coughing/ Dry mouth/ Nausea/ Feeling drunk/ Sweating. Uncommon: Tingling or numbness to hands and feet/ Tiredness/ Mouth discomfort.
Additional information	Patients with pain due to acute coronary syndrome (ACS) or migraine may not be suitable for Methoxyflurane. Methoxyflurane crosses the placenta. Consider the risk of central nervous system (CNS) and respiratory depression in an already compromised foetus. Methoxyflurane has a mildly pungent odour. If used in a confined space request the patient to inhale and exhale through the inhaler tube while ensuring that the activated Carbon Chamber is attached.

Clinical Level:



MEDICATION	NALOXONE
Classification	Opioid toxicity: Opioid receptor antagonist. The management and reversal of opiate overdose.
Presentation	Ampoules 400 mcg/mL (0.4 mg in 1 mL) / Minijet syringe.
Administration	IV / IO / IM / SC / IN. (CPG: 6.10.2, 4/5/6.12.7, 4/5/6.13.7 4/5/6.14.6).
Indications	Inadequate respiration and/or ALoC following known or suspected narcotic overdose.
Contra-Indications	Known severe adverse reaction.
Usual Dosages	<p>Adult: 400 mcg IV/IO (AP) (repeat after 3 min PRN to a Max dose of 2 mg). 400 mcg IM/SC (P) (repeat after 3 min PRN to a Max dose of 2 mg). 800 mcg IN (EMT) (repeat x 1 after 3 min PRN).</p> <p>Paediatric: 10 mcg/kg IV/IO (AP). 10 mcg/kg IM/SC (P). 20 mcg/kg IN (EMT). (Repeat dose PRN to maintain opioid reversal to Max 0.1 mg/kg or 2 mg).</p>
Side effects	Acute reversal of narcotic effect ranging from nausea and vomiting to agitation and seizures.
Additional information	Use with caution in pregnancy. Administer with caution to patients who have taken large dose of narcotics or are physically dependent. Rapid reversal will precipitate acute withdrawal syndrome. Prepare to deal with aggressive patients.

Clinical Level:



MEDICATION	NITROUS OXIDE 50% AND OXYGEN 50% (ENTONOX®)
Classification	Analgesics – Volatile Liquid Anaesthetics - Potent analgesic gas contains a mixture of both Nitrous Oxide and Oxygen.
Presentation	Cylinder, coloured blue with white and blue triangles on cylinder shoulders. ED cylinder: White cylinder. <i>Medical gas:</i> 50% Nitrous Oxide & 50% Oxygen. Brand name: Entonox®.
Administration	Self-administered. Inhalation by demand valve with face-mask or mouthpiece. (CPG: 4/5/6.6.2, 4/5/6.12.3, 4/5/6.12.4, 4/5/6.13.13).
Indications	Moderate to severe pain.
Contra-Indications	Altered level of consciousness/ Chest Injury/ Pneumothorax/ Shock / Recent scuba dive/ Decompression sickness/ Intestinal obstruction/ Inhalation Injury/ Carbon monoxide (CO) poisoning/ Known severe adverse reaction.
Usual Dosages	<i>Adult and Paediatric:</i> Self-administered until pain tolerable.
Side effects	Disinhibition/ Decreased level of consciousness/ Light headedness.
Additional information	Caution should be issued before using Entonox with patients who have known Chronic Obstructive Pulmonary Disease (COPD) or other conditions where compromised chemoreceptor sensitivity/function may be present. May cause respiratory depression and increases in PaCO ₂ . Do not use if patient unable to understand instructions. In cold temperatures warm cylinder and invert at least 3 times to ensure mix of gases. Advanced paramedics may use discretion with minor chest injuries. Has an addictive property. Caution when using Entonox® for greater than one hour for sickle cell crisis. Prolonged or frequent use of ENTONOX may result in megaloblastic marrow changes, myeloneuropathy and sub-acute combined degeneration of the spinal cord.

Clinical Level:



MEDICATION	OXYGEN
Classification	Gas.
Presentation	<p><i>Medical gas:</i> D, E or F cylinders, coloured black with white shoulders. (Please note: By 2025, all cylinders will be completely white with OXYGEN in black). <i>CD cylinder:</i> White cylinder.</p>
Administration	<p><i>Inhalation via:</i> High concentration reservoir (non-rebreather) mask/ Simple face mask/ Venturi mask/ Tracheostomy mask/ Nasal cannulae/ CPAP device/ Bag Valve Mask. (CPG: Oxygen is used extensively throughout the CPGs).</p>
Indications	<p>Absent / Inadequate ventilation following an acute medical or traumatic event. SpO₂ < 94% adults and < 96% paediatrics. SpO₂ < 92% for patients with acute exacerbation of COPD. SpO₂ < 90% for patients with acute onset of Pulmonary Oedema.</p>
Contra-Indications	Bleomycin lung injury.
Usual Dosages	<p>Adult: Cardiac and respiratory arrest or sickle cell crisis; 100%. Life threats identified during primary survey; 100% until a reliable SpO₂ measurement obtained then titrate O₂ to achieve SpO₂ of 94% - 98%. For patients with acute exacerbation of COPD, administer O₂ titrate to achieve SpO₂ 92% or as specified on COPD Oxygen Alert Card. All other acute medical and trauma titrate O₂ to achieve SpO₂ 94% - 98%.</p> <p>Paediatric: Cardiac and respiratory arrest or sickle cell crisis; 100%. Life threats identified during primary survey; 100% until a reliable SpO₂ measurement obtained then titrate O₂ to achieve SpO₂ of 96% - 98%. Neonatal resuscitation (< 4 weeks) consider supplemental O₂ (≤ 30%). All other acute medical and trauma titrate O₂ to achieve SpO₂ of 96% - 98%.</p>
Side effects	Prolonged use of O ₂ with chronic COPD patients may lead to reduction in ventilation stimulus.
Additional information	<p>Caution with emollients containing paraffin e.g. lip balms & moisurisers – may lead to skin burns. A written record must be made of what oxygen therapy is given to every patient. Documentation recording oximetry measurements should state whether the patient is breathing air or a specified dose of supplemental Oxygen. Consider humidifier if oxygen therapy for paediatric patients is > 30 minutes duration. Caution with paraquat poisoning, administer Oxygen if SpO₂ < 92%. Avoid naked flames, powerful oxidising agent.</p>

Clinical Level:



MEDICATION	PARACETAMOL
Classification	Analgesic – Non-opioid.
Presentation	Rectal suppository 1 g, 500 mg, 250 mg, 180 mg, 125 mg, 80 mg. Suspension 120 mg in 5 mL or 250 mg in 5 mL. 500 mg tablet. Plastic vial, 1 g of Paracetamol in 100 mL solution for infusion, 500mg of paracetamol in 50 mL solution for infusion.
Administration	Per Rectum (PR). Orally (PO). IV infusion. (CPG: 4/5/6.6.2, 4/5/6.11.1, 4/5/6.13.13, 4/5/6.13.19, 5/6.13.20, 5/6.15.2).
Indications	<i>Adult:</i> Pyrexia/ Temperature > 38.3°C/ Mild or moderate pain. <i>Paediatric:</i> Pyrexia/ Temperature > 38.5°C/ Mild or moderate pain.
Contra-Indications	< 1 month old/ Known severe adverse reaction/ Chronic liver disease.
Usual Dosages	<i>Adult:</i> 1 g PO (EMT, P/AP). 1 g IV infusion (AP), if estimated weight < 50 kg, 15 mg/kg (administered slowly over 15 minutes). <i>Palliative Care:</i> 1g PO (Repeat after 4-6 hours x 1 PRN). <i>Paediatric:</i> PO (EMT, P/AP) PR (AP) IV Infusion (AP) (≥ 1 year Max 1g) 15 mg/Kg PO > 1 month < 1 year - 80 mg PR < 1 year – 7.5 mg/kg IV slowly 1-3 years - 180 mg PR ≥ 1 year – 15 mg/kg IV slowly 4-8 years - 360 mg
Side effects	If Paracetamol IV is administered too fast it may result in hypotension.
Additional information	Paracetamol is contained in Paracetamol suspension and other over the counter drugs. Consult with parent / guardian in relation to medication administration prior to arrival on scene. For PR use be aware of the modesty of the patient, should be administered in the presence of a 2nd person. If Paracetamol administered in the previous 4 hours, adjust the dose downward by the amount given by other sources resulting in a maximum of 15 mg/Kg. Caution with IV Paracetamol in the absence of a buretrol.

Clinical Level:



MEDICATION	SALBUTAMOL
Classification	Beta-2 Adrenoceptor agonist selective – short acting.
Presentation	Nebule 2.5 mg in 2.5 mL. Nebule 5 mg in 2.5 mL. Aerosol inhaler: Metered dose 100mcg per actuation (Puff).
Administration	Nebule Inhalation via aerosol inhaler. (CPG: 4/5/6.2.4, 2/3.2.5, 4/5/6.2.5, 4/5/6.8.9, 2/3.10.1, 4/5/6.10.1, 2/3.13.8, 4/5/6.13.8, 2/3.13.21, 4/5/6.13.21, 6.17.7).
Indications	Bronchospasm/ Exacerbation of COPD/ Respiratory distress following submersion incident.
Contra-Indications	Known severe adverse reaction.
Usual Dosages	Adult: 5 mg NEB or 100mcg metered aerosol spray (repeat aerosol x 11). Repeat NEB at 5 minute intervals PRN EFR assist patient with Asthma/ Anaphylaxis. - 100mcg metered aerosol spray (repeat aerosol x 11 PRN). Paediatric: < 5 yrs - 2.5 mg NEB or 100 mcg metered aerosol spray (repeat aerosol x 5). > 5 yrs - 5 mg NEB or 100 mcg metered aerosol spray (repeat aerosol x 11). (Repeat NEB at 5 minute intervals PRN). EFR: assist patient with Asthma/ Anaphylaxis – < 5 yrs - 100 mcg/ 1 actuation metered aerosol spray (repeat aerosol x 5 PRN). > 5 yrs - 100 mcg/ 1 actuation metered aerosol spray (repeat aerosol x 11 PRN).
Side effects	Tachycardia/ Tremors/ Tachyarrhythmias/ High doses may cause Hypokalaemia.
Additional information	It is more efficient to use a volumiser in conjunction with an aerosol inhaler when administering Salbutamol. If an oxygen driven nebuliser is used to administer Salbutamol for a patient with acute exacerbation of COPD it should be limited to 6 minutes maximum.

New Medications and Skills for 2021

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Activated Charcoal PO*					√	√	√
Adrenaline nebulised						√	√
Dexamethasone PO/IM						√	√
Lidocaine IO							√
Ketamine IM*							√
Uterine massage					√	√	√
Tourniquet application					√	√	√
Pressure points					√	√	√
Ketone measurement*					√	√	√
Tracheostomy management					√	√	√
Malpresentations in labour						√	√
Shoulder Dystocia management						√	√
Posterior ECG in ACS						√	√
Intubation of Stoma							√
Nasogastric Tube insertion*							√
Procedural Sedation*							√
Richmond Agitation-Sedation Scale (RASS)*							√

Care management including the administration of medications as per level of training and division on the PHECC Register and Responder levels.

Pre-Hospital Responders and Practitioners shall only provide care management including medication administration for which they have received specific training. Practitioners must be privileged by a licensed CPG provider to administer specific medications and perform specific clinical interventions.

√	Authorised under PHECC CPGs
URMPIO	Authorised under PHECC CPGs under registered medical practitioner's instructions only
APO	Authorised under PHECC CPGs to assist practitioners only (when applied to EMT to assist paramedic or higher clinical levels)
√ SA	Authorised subject to special authorisation as per CPG
BTEC	Authorised subject to Basic Tactical Emergency Care rules
*	Non-core specified element or action
√*	Non-core specified element or action for identified clinical level

Paramedic authorisation for IV continuation

Practitioners should note that PHECC registered paramedics are authorised to continue an established IV infusion in the absence of an advanced paramedic or doctor during transportation.

MEDICATIONS

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Aspirin PO	√	√	√	√	√	√	√
Oxygen INH		√		√	√	√	√
Glucose gel buccal				√	√	√	√
Glyceryl Trinitrate SL				√ SA	√	√	√
Adrenaline (1:1000) autoinjector				√ SA	√	√	√
Salbutamol MDI				√ SA	√	√	√
Activated Charcoal PO*					√	√	√
Adrenaline (1:1000) IM					√	√	√
Chlorphenamine PO/IM					√	√	√
Glucagon IM					√	√	√
Ibuprofen PO					√	√	√
Methoxyflurane INH					√	√	√
Naloxone IN					√	√	√
Nitrous Oxide and Oxygen INH					√	√	√
Paracetamol PO					√	√	√
Salbutamol nebulised					√	√	√
Adrenaline nebulised						√	√
Clopidogrel PO						√	√
Cyclizine IM						√	√
Dexamethasone PO/IM						√	√
Glucose 5% IV						√ SA	√
Glucose 10% IV						√ SA	√
Hydrocortisone IM						√	√
Ipratropium Bromide nebulised						√	√
Midazolam buccal/IM/IN						√	√

APPENDIX 2 - Medication & Skills Matrix

EMERGENCY MEDICAL TECHNICIAN

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Naloxone IM/SC						√	√
Ondansetron IM						√	√
Oxytocin IM						√	√
Ticagrelor PO						√	√
Sodium Chloride 0.9% IV/IO						√ SA	√
Adenosine IV							√
Adrenaline (1:10,000) IV/IO							√
Amiodarone IV/IO							√
Atropine IV/IO							√
Ceftriaxone IV/IO/IM							√
Chlorphenamine IV							√
Cyclizine IV							√
Diazepam IV/PR							√
Fentanyl IN/IV							√
Furosemide IV							√
Glycopyrronium Bromide SC*							√
Haloperidol PO/SC*							√
Hydrocortisone IV							√
Hyoscine Butylbromide SC*							√
Ketamine IV/IM*							√
Lidocaine IV/IO							√
Lorazepam PO							√
Magnesium Sulphate IV							√
Midazolam IV							√
Morphine IV/PO/IM							√
Naloxone IV/IO							√
Ondansetron IV							√
Paracetamol IV/PR							√
Sodium Bicarbonate IV/IO							√
Tranexamic Acid IV							√

AIRWAY & BREATHING MANAGEMENT

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
FBAO management	✓	✓	✓	✓	✓	✓	✓
Head tilt chin lift	✓	✓	✓	✓	✓	✓	✓
Pocket mask	✓	✓	✓	✓	✓	✓	✓
Recovery position	✓	✓	✓	✓	✓	✓	✓
Non-rebreather mask		✓		✓	✓	✓	✓
Oropharyngeal airway		✓		✓	✓	✓	✓
Oral suctioning		✓		✓	✓	✓	✓
Venturi mask		✓		✓	✓	✓	✓
Bag Valve Mask		✓		✓	✓	✓	✓
Jaw thrust				✓	✓	✓	✓
Nasal cannula		✓		✓	✓	✓	✓
Oxygen humidification				✓	✓	✓	✓
Nasopharyngeal airway				BTEC	BTEC	✓	✓
Supraglottic airway adult (uncuffed)		✓			✓	✓	✓
Supraglottic airway adult (cuffed)					✓ SA	✓	✓
Tracheostomy management					✓	✓	✓
Continuous Positive Airway Pressure						✓	✓
Non-Invasive ventilation device						✓	✓
Supraglottic airway paediatric						✓	✓
Endotracheal intubation							✓
Intubation of stoma							✓
Laryngoscopy / Magill forceps							✓
Needle cricothyrotomy							✓
Needle thoracocentesis							✓

CARDIAC

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
AED adult & paediatric	√	√	√	√	√	√	√
CPR adult, child & infant	√	√	√	√	√	√	√
Recognise death and resuscitation not indicated	√	√	√	√	√	√	√
Neonate resuscitation					√	√	√
ECG monitoring					√	√	√
CPR mechanical assist device*					√	√	√
Cease resuscitation - adult					√ SA	√	√
12 lead ECG						√	√
Manual defibrillation						√ *	√
Right sided ECG in ACS						√	√
Posterior ECG in ACS						√	√

HAEMORRHAGE CONTROL

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Direct pressure			√	√	√	√	√
Nose bleed			√	√	√	√	√
Haemostatic agent				BTEC*	√ *	√	√
Tourniquet application				BTEC	√	√	√
Pressure points					√	√	√
Wound closure clips					BTEC	√ *	√ *
Nasal pack						√	√

MEDICATION ADMINISTRATION

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Oral	✓	✓	✓	✓	✓	✓	✓
Buccal				✓	✓	✓	✓
Metered dose inhaler				✓ SA	✓	✓	✓
Sublingual				✓ SA	✓	✓	✓
Intramuscular injection					✓	✓	✓
Intranasal					✓	✓	✓
Nebuliser					✓	✓	✓
Subcutaneous injection					✓	✓	✓
Infusion maintenance						✓	✓
Infusion calculations							✓
Intraosseous injection/infusion							✓
Intravenous injection/infusion							✓
Per rectum							✓

TRAUMA

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Burns care			✓	✓	✓	✓	✓
Application of a sling			✓	✓	✓	✓	✓
Soft tissue injury			✓	✓	✓	✓	✓
Active Spinal Motion Restriction			✓	✓	✓	✓	✓
Hot packs for active rewarming (hypothermia)			✓	✓	✓	✓	✓
Cervical collar application				✓	✓	✓	✓
Helmet stabilisation/removal				✓	✓	✓	✓
Splinting device application to upper limb				✓	✓	✓	✓
Splinting device application to lower limb				✓	✓	✓	✓
Log roll				APO	✓	✓	✓
Move patient with a carrying sheet				APO	✓	✓	✓
Extrication using a long board				✓ SA	✓	✓	✓
Rapid Extraction				✓ SA	✓	✓	✓
Secure and move a patient with an extrication device				✓ SA	✓	✓	✓
Move a patient with a split device (Orthopaedic stretcher)				✓ SA	✓	✓	✓
Passive Spinal Motion Restriction						✓	✓
Pelvic Splinting device				BTEC	✓	✓	✓
Move and secure patient into a vacuum mattress				BTEC	✓	✓	✓
Move and secure a patient to a paediatric board					✓	✓	✓
Traction splint application					APO	✓	✓
Lateral dislocation of patella – reduction						✓	✓
Taser gun barb removal						✓	✓

OTHER

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Use of Red Card	√	√	√	√	√	√	√
Assist normal delivery of a baby				APO	√	√	√
De-escalation and breakaway skills					√	√	√
ASHICE radio report					√	√	√
IMIST-AMBO handover					√	√	√
Uterine massage					√	√	√
Malpresentations in labour						√	√
Shoulder Dystocia management						√	√
Umbilical cord complications						√	√
Verification of Death						√	√
Intraosseous cannulation							√
Intravenous cannulation							√
Nasogastric tube insertion*							√
Procedural Sedation*							√
Urinary catheterisation*							√

PATIENT ASSESSMENT

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
Assess responsiveness	√	√	√	√	√	√	√
Check breathing	√	√	√	√	√	√	√
FAST assessment	√	√	√	√	√	√	√
Capillary refill			√	√	√	√	√
AVPU			√	√	√	√	√
Pulse check			√	√	√	√	√
Breathing / pulse rate		√ SA	√	√	√	√	√
Primary survey			√	√	√	√	√
SAMPLE history			√	√	√	√	√
Secondary survey			√	√	√	√	√

APPENDIX 2 - Medication & Skills Matrix

EMERGENCY MEDICAL TECHNICIAN

CLINICAL LEVEL	CFR-C	CFR-A	FAR	EFR	EMT	P	AP
CSM assessment				√	√	√	√
Rule of Nines				√	√	√	√
Assess pupils				√	√	√	√
Blood pressure				√ SA	√	√	√
Capacity evaluation					√	√	√
Chest auscultation					√	√	√
Glucometry					√	√	√
Ketone measurement*					√	√	√
Paediatric Assessment Triangle					√	√	√
Pain assessment					√	√	√
Patient Clinical Status					√	√	√
Pulse oximetry					√	√	√
Temperature					√	√	√
Triage sieve					√	√	√
Broselow tape						√	√
Capnography						√	√
Glasgow Coma Scale (GCS)						√	√
Peak expiratory flow						√	√
Pre-hospital Early Warning Score						√	√
Treat and referral						√	√
Triage sort						√	√
Richmond Agitation-Sedation Scale (RASS) *							√

CRITICAL INCIDENT STRESS MANAGEMENT (CISM)

Your Psychological Well-Being

It is extremely important for your psychological well-being that you do not expect to save every critically ill or injured patient that you treat. For a patient who is not in hospital, whether they survive a cardiac arrest or multiple traumas depends on a number of factors including any other medical condition the patient has. Your aim should be to perform your interventions well and to administer the appropriate medications within your scope of practice. However, sometimes you may encounter a situation which is highly stressful for you, giving rise to Critical Incident Stress (CIS). A critical incident is an incident or event which may overwhelm or threaten to overwhelm our normal coping responses. As a result of this we can experience CIS.

When can I be adversely affected by a critical incident? Listed below are some common ways in which people react to incidents like this:

- Feeling of distress or sadness
- Strong feeling of anger
- Feeling of disillusionment
- Feeling of guilt
- Feeling of apprehension/anxiety/fear of:
 - Losing control/breaking down or
 - Something similar happening again
 - Not having done all I think I could have done
- Avoidance of the scene of incident/trauma
- Bad dreams, nightmares or startling easily
- Distressing memories or 'flashbacks' of the incident
- Feeling 'on edge', irritable, angry, under threat/pressure
- Feeling emotionally fragile or emotionally numb
- Feeling cut off from your family or close friends – "I can't talk to them" or "I don't want to upset them"
- Feeling of needing to control everything

Some Do's and Don'ts

- **DO** express your emotions:
 - Talk about what happened
 - Talk about how you feel and how the event has impacted you
 - Be kind to yourself and to others.
- **DO** talk about what has happened as often as you need
- **DO** find opportunities to review the experience **DO** discuss what happened with colleagues **DO** ask friends and colleagues for support
- **DO** listen sympathetically if a colleague wants to talk
- **DO** advise colleagues about receiving appropriate help
- **DO** keep to daily routines
- **DO** drive more carefully
- **DO** be more careful around the home
- **DON'T** use alcohol, nicotine or drugs to hide your feelings **DON'T** simply stay away from work – seek help and support **DON'T** allow anger and irritability to mask your feelings **DON'T** bottle up feelings
- **DON'T** be afraid to ask for help
- **DON'T** think your feelings are a sign of weakness

When things get tough, pro-actively minding yourself is crucial. Control the things you can control. Get more sleep than you think you need. Eat fresh, healthy foods at regular times and avoid snacks. Get outdoor exercise at least three times a week. Have a meaningful conversation with someone you like at least once a day. Resolve what makes you sad or angry or otherwise let it go. Be kind.

Everyone may have these feelings. Experience has shown that they may vary in intensity according to circumstance. Nature heals through allowing these feelings to come out. This will not lead to loss of control but stopping these feelings may lead to other and possibly more complicated problems.

When to find help?

1. If you feel you cannot cope with your reactions or feelings.
2. If your stress reactions do not lessen in the two or three weeks following the event.
3. If you continue to have nightmares and poor sleep.
4. If you have no-one with whom to share your feelings when you want to do so.
5. If your relationships seem to be suffering badly, or sexual problems develop.
6. If you become clumsy or accident prone.
7. If, in order to cope after the event, you smoke, drink or take more medication, or other drugs.
8. If your work performance suffers.
9. If you are tired all the time.
10. If things get on top of you and you feel like giving up.
11. If you take it out on your family.
12. If your health deteriorates.

Experiencing signs of excessive stress?

If the range of physical, emotional and behavioural signs and symptoms already mentioned do not reduce over time (for example after two weeks), it is important that you seek support and help.

Where to find help?

Your own licensed CPGs provider will have a CISM support network or system.

We recommend that you contact them for help and advice (i.e. your peer support worker/ coordinator/staff support officer).

- For a self-help guide, please go to www.cismnetworkireland.ie
- The NAS CISM and CISM Network published a booklet called 'Critical Incident Stress Management for Emergency Personnel'.
- It can be purchased by emailing: info@cismnetworkireland.ie
- Consult your own GP or see a health professional who specialises in traumatic stress.
- In partnership with NAS CISM Committee, PHECC developed an eLearning CISM Stress Awareness Training (SAT) module. It can be accessed by the following personnel:
 - PHECC registered practitioners at all levels
 - National Ambulance Service-linked community first responders
 - NAS non-PHECC registered personnel
- Under the direction of CISM Network, bespoke CISM SAT modules are developed by Network member organisations.

Practitioner Level Updates

Several broad changes have been applied in the 2021 version:

- Care Principles have been updated.
- The classification of CPGs has changed to up to seventeen categories, developed to group common themes and categories together.
- The term 'Registered' has been removed from references to registered healthcare professionals, for example Registered Medical Practitioner will now appear as Medical Practitioner.
- The transport patient symbol, along with other symbols, has been modernised throughout the CPGs.
- The description of dose of medications less than one milligram is now described in micrograms, for example GTN 0.4 mg SL is now GTN 400 mcg SL.
- The description of sodium chloride (0.9%) infusion has been standardised to NaCl (0.9%).
- Epinephrine is now known as Adrenaline throughout the CPGs.
- The Medical Support symbol now states 'Consider Medical Support or 'Contact Medical Support'. Where 'Contact Medical Support' appears this should be regarded as mandatory.
- References to published source literature no longer appear on CPGs but are available from PHECC on request.
- The age descriptor has been removed from the title of paediatric CPGs.

New EMT CPGs in 2021 Edition

To support upskilling of the 2021 CPGs new CPGs are identified below.

New CPGs	The new skills and medications incorporated into the CPGs are
4/5/6.2.7 Emergency Tracheostomy Management	This CPG outlines the approach to managing respiratory issues in a patient with a tracheostomy. Includes saline nebulised
CPG 4/5/6.12.1 Pregnancy related emergencies	This CPG outlines the assessment and management of pregnancy related emergencies.

New CPGs	The new skills and medications incorporated into the CPGs are
CPG 4/5/6.12.2 Pre-Hospital Emergency Childbirth	This CPG replaces the previous Pre-Hospital Emergency Childbirth CPG and integrates EMT, Paramedic and Advanced Paramedic.
CPG 4/5/6.12.7 New-born Neonatal Care and Resuscitation	This CPG outlines the assessment and management of the new-born including resuscitation.
CPG 4/5/6.12.8 Neonatal Resuscitation (≤ 6 weeks)	This CPG outlines the approach to neonatal resuscitation and integrates EMT, Paramedic and Advanced Paramedic.
CPG 4/5/6.13.18 Limb Injury – Paediatric	This CPG outlines the approach to paediatric limb injury.
CPG 4/5/6.12.20 Sepsis – Paediatric	This CPG outlines the approach to paediatric sepsis.

Deleted EMT CPGs in 2021 Edition

CPGs deleted	
CPG 4.5.1 Pre-Hospital Emergency Childbirth	This CPG has been deleted and replaced with CPG 4/5/6.12.2 Pre-Hospital Emergency Childbirth which integrates EMT, Paramedic and Advanced Paramedic levels.
CPG 4/5/6.5.3 PV Haemorrhage in Pregnancy	This CPG has been deleted and replaced with CPG 4/5/6.12.1 – Pregnancy related emergencies.
CPG 4.5.2 Basic Life Support – Neonate (< 4 week)	This CPG has been deleted and replaced with CPG 4/5/6.12.8 Neonatal Resuscitation (≤ 6 weeks) which integrates EMT, Paramedic and Advanced Paramedic levels.

Updated EMT CPGs in the 2021 version

To support upskilling of the 2021 CPGs, the changes are outlined below.

CPGs	The principal differences are
CPG 4/5/6.2.3 Abnormal Work of Breathing – Adult	The CPG is retitled 'Abnormal Work of Breathing – Adult' (previously Inadequate Ventilations – Adult)

CPGs	The principal differences are
CPG 4/5/6.2.4 Exacerbation of COPD	<p><i>Deleted</i></p> <p>Sequence step 'Measure Peak Expiratory Flow'</p> <p>Decision process 'PEF < 50% predicted'</p> <p><i>Added</i></p> <p>Decision process 'Deteriorates / unstable' replaces decision process 'PEF < 50% predicted'</p> <p>Decision process 'Adequate ventilation' replaces decision process 'Adequate respirations'</p> <p>Consider treatment 'consider CPAP for profound refractory hypoxia' is introduced for Paramedic and AP level</p> <p>Instruction box 'If no improvement Salbutamol may be repeated at 5 min intervals'</p> <p><i>Medication Update</i></p> <p>Salbutamol may be repeated at 5 min intervals</p>
CPG 4/5/6.2.5 Asthma - Adult	<p><i>Added</i></p> <p>Instruction box 'If no improvement Salbutamol aerosol 100mcg may be repeated up to 11 times as required <u>via MDI</u>' replaces 'If no improvement Salbutamol aerosol 0.1 mg may be repeated up to 11 times as required'</p>
CPG 4.3.1 Acute Coronary Syndrome	<p><i>Added</i></p> <p>Parallel process 'Contact Ambulance Control/ Operations Centre for direction' replaces 'Contact NAS control for direction'</p>
4/5/6.4.2 Epistaxis	<p><i>Deleted</i></p> <p>Equipment list 'Proprietary nasal pack'</p>
CPG 4/5/6.5.2 Decompression Illness	<p><i>Added</i></p> <p>Transport patient 'Transport is completed at an altitude of < 1000 ft. above incident site or aircraft pressurised equivalent to sea level' replaces 'Transport is completed at an altitude of < 300 meters above incident site or aircraft pressurised equivalent to sea level'</p>
CPG 4/5/6.5.3 Glycaemic Emergency – Adult	<p><i>Added</i></p> <p>Consider treatment option 'Consider Ketone measurement' is a non-core element for EMT, Paramedic and AP level</p> <p>Instruction box 'Consider Glucagon IM if not already given'</p>
CPG 4/5/6.5.4 Sickle Cell Crisis - Adult	<p><i>Added</i></p> <p>Instruction box 'Administer 15L of oxygen via a non-rebreather facemask' replaces '100% O₂'</p>

CPGs	The principal differences are
CPG 4.6.1 Altered Level of Consciousness - Adult	<i>Added</i> 'Possible differential diagnosis' box replaces 'Differential diagnosis' box
CPG 4/5/6.6.2 Pain Management – Adult	<i>Deleted</i> Instruction box 'Following Fentanyl IN the next dose may be either Fentanyl IV or Morphine IV but not both' <i>Added</i> Instruction box 'Repeat Fentanyl IN once only at not < 10 min after initial dose prn' Instruction box 'Repeat Ketamine PRN at not < 10 min' replaces 'Repeat Ketamine once only at not < 10 minutes prn' Instruction box 'Poly-opiate administration should be avoided where possible – where multiple opiates are being administered the highest standards of continued patient monitoring must be adhered to' Cyclical process box for 'IO Access & Analgesia' Special instructions box 'Do not administer Amiodarone and Lidocaine to the same patient'
CPG 4/5/6.7.2 Behavioral Emergency	<i>Deleted</i> 'or if the implementation of the decision requires the act of a third party' from the Instruction box describing the circumstances where a person lacks the capacity to make a decision <i>Added</i> Decision process 'Aggressive/violent and/or risk to self or others and uncooperative with practitioner' is reorganised Mandatory sequence step 'Hand over to MP/ Garda care' replaces 'Hand over to RMP/ Garda care' Sequence step 'ETCO ₂ ' added to 'Monitor BP SpO ₂ and ECG' Sequence step 'Mental Health Illness' Instruction box 'If potential to harm self or others ensure minimum two people accompany patient in saloon of ambulance at all times' replaces 'Consider need for two or more people accompanying the patient during transportation'

CPGs	The principal differences are
<p>CPG 4/5/6.8.3 External Haemorrhage – Adult</p>	<p><i>Deleted</i></p> <p>Paramedic skill flag from mandatory sequence step 'Apply tourniquet if limb injury'</p> <p>Paramedic skill flag from sequence step 'Depress proximal pressure point'</p> <p>Paramedic skill flag from sequence step 'Apply tourniquet'</p> <p>'apply a tourniquet and/or' from EMT-BTEC Special Authorisation box</p> <p><i>Added</i></p> <p>Mandatory sequence step 'Apply and mark tourniquet if limb injury' replaces 'Apply tourniquet if limb injury' and is introduced as an EMT level skill</p> <p>Consider treatment option 'Consider wound closure clips for temporary closure if serious haemorrhage' is a non-core element for Paramedic and AP level</p> <p>Consider treatment option 'consider applying a dressing impregnated with haemostatic agent' is a Paramedic level skill</p> <p>Consider treatment option 'consider applying a dressing impregnated with haemostatic agent' is a non-core element for EMT level</p>
<p>CPG 4/5/6.8.6 Limb Injury – Adult</p>	<p>The CPG treatment pathway is reorganised</p> <p><i>Deleted</i></p> <p>Instruction box 'Contraindications for application of traction splint'</p> <p>Decision process 'Injury type'</p> <p>Decision process '> 20 min to facility'</p> <p><i>Added</i></p> <p>Parallel process 'Consider hypovolaemia and shock'</p> <p>Decision process 'Fracture'</p> <p>Decision process 'Femur fracture' replaces clinical finding 'Fractured femur'</p> <p>Decision process 'Rest, Cooling, Compression & Elevation' replaces decision process Rest – Ice – Compression – Elevation'</p> <p>Sequence step 'For open fracture - Remove gross contamination'</p>

CPGs	The principal differences are
<p>CPG 4/5/6.8.9 Submersion / Immersion Incident</p>	<p>The CPG is retitled 'Submersion/ Immersion Incident' (previously Submersion Incident) The CPG entry point is updated to 'Submersion / immersion in liquid' The CPG treatment pathway is reorganised Instruction box outlining the indications of spinal injury is revised to 'History of: Diving into shallow water – Injury following: water slide, water skiing, kite-surfing, boat incident – Alcohol/ drugs intoxication'</p> <p><i>Deleted</i> Decision process 'Adequate ventilations'</p> <p><i>Added</i> Decision process 'Responsive' Decision process 'Spontaneous Breathing' Mandatory sequence step 'Open airway - Five rescue breaths' Decision process 'Spontaneous Breathing' Mandatory sequence step 'Continue ventilations' Sequence step 'Advanced airway with cuffed devices only (monitor for leaks)' for Paramedic and AP level Consider treatment option 'Consider nasogastric tube' is a non-core element for AP level Sequence step 'Auscultate lungs' Decision process 'Crepitations' Decision process 'Hypotensive' Decision process Hypothermic' replaces 'Patient is hypothermic'</p>
<p>CPG 4/5.10.2 Poisons – Adult</p>	<p><i>Added</i> Additional CPG entry point 'Solid substance ingested and GCS 15' Decision process 'Activated charcoal indicated' Instruction box 'Substances that are adsorbed by Activated charcoal are available in the PHECC field guide'.</p> <p>Medication Update 'Naloxone 0.8 - 2 mg IN (Repeat x one prn to max 2mg)' replaces 'Naloxone 0.8 mg IN (Repeat x one prn)'</p> <p>New Medications Consider treatment option 'Activated charcoal 50 g PO' is a non-core element for Paramedic and EMT level</p>

CPGs	The principal differences are
<p>CPG 4/5/6.11.1 Sepsis - Adult</p>	<p>The CPG entry point is updated to 'Patient generally unwell with suspected infection < 36°C or > 38.3°C'</p> <p>The CPG treatment pathway is significantly reorganised</p> <p><i>Deleted</i></p> <p>Sequence step 'Signs of Systemic Inflammatory Response Syndrome (SIRS)'</p> <p>Sequence step 'Could this be a severe infection?'</p> <p>Instruction box 'Risk stratifier'</p> <p>Instruction box 'If history of penicillin allergy assess the severity of the reaction and if not life-threatening, i.e. rash, proceed with Ceftriaxone'</p> <p>Instruction box 'If meningitis suspected ensure appropriate PPE is work; Mask and goggles'</p> <p>Instruction box 'Indication for antibiotic'</p> <p>Instruction box 'Signs of shock/ poor perfusion'</p> <p>Consider treatment 'If Sys BP < 100 mmHg consider aliquots NaCl 0.9% 250 mL IV/IO</p> <p><i>Added</i></p> <p>Sequence step 'HR, RR, ECG, SpO₂ & BP monitoring' replaces 'ECG, SpO₂ & BP monitoring'</p> <p>Mandatory sequence step 'Abnormal physiology? Source of Infection?'</p> <p>Decision process 'At risk'</p> <p>Instruction box 'Evidence of at risk criteria (any 1 of 3) 1. Any 1 Clinical sign of acute organ dysfunction - 2. At risk of neutropenia (bone marrow failure, autoimmune disorder, treatment including but not limited to chemo/ radiotherapy). Note: these patients may present without fever - 3. ≥ SIRS criteria PLUS ≥ 1 co-morbidity'</p> <p>Sequence step 'Give 3 if clinically indicated'</p> <p>Instruction box 'Signs of Systemic Inflammatory Response Syndrome (SIRS)'</p> <p>Instruction box 'Give 3 1. O₂ titrate to sats > 94% (88 – 92% for chronic lung conditions e.g. COPD) 2. IV fluids, patients with hypotension max 30 mL/kg 3. IV antimicrobials' replaces 'Give three - O₂ titrate to sats > 94% - IV fluids – IV antimicrobials'</p> <p>Decision process 'Signs of hypoperfusion' replaces 'Signs of poor perfusion'</p> <p>Sequence step 'Monitor clinical condition; re-evaluate for possible sepsis if clinically indicated'</p> <p>Instruction box 'High Consequence Infectious Disease (HCID) ensure appropriate PPE is worn; Long sleeve gown, Facemask, Eye protection'</p> <p>Special instruction box 'Pre alert ED if sepsis" replaces 'Pre alert ED if severe sepsis'</p>

CPGs	The principal differences are
<p>CPG 4/5/6.12.2 Pre-Hospital Emergency Childbirth</p>	<p>EMT level is added to this CPG The CPG treatment pathway is significantly reorganised The CPG entry point is updated to 'In labour'</p> <p><i>Deleted</i> Instruction box 'If no progress with labour consider transporting patient' Sequence step 'Take SAMPLE history' Decision process 'Patient in labour' Mandatory sequence step 'Request Ambulance Control to contact GP/ midwife/medical team as required by local policy to come to scene or meet en route' Mandatory sequence step 'Cover newborn in polythene wrap/bag up to neck without drying first'</p> <p><i>Added</i> Decision process 'Birth imminent' replaces 'Birth imminent or travel time too long' 'Consider ALS' replaces 'Request ALS' Sequence step 'Request second crew' Instruction box 'Consider Additional crew for each baby expected' Decision process 'Malpresentation' replaces 'Breech birth' Decision process 'Risk factors' Sequence step 'Initiate rapid transport – Pre-alert labour ward – Optimise resuscitation of mother' Mandatory sequence step 'Aim: birth in hospital' Sequence step 'Mother to adopt position of comfort and prepare environment & equipment for birth' replaces 'Position mother and prepare equipment for birth' Sequence step 'Monitor vital signs' replaces 'Monitor vital signs and BP' Decision process 'Pre-hospital delivery' Mandatory sequence step 'Warm, dry, stimulate baby. Check ABCs' replaces 'Dry baby and check ABCs' Mandatory sequence step 'Check for second baby' Sequence step 'Skin to skin contact' replaces 'Wrap baby and present to mother (Skin to skin preference)' Sequence step 'Encourage breastfeeding (no contraindications)' Instruction box 'Risk factors for complicated delivery – Prematurity – Multiple births – PV bleeding – Pre-eclampsia indicators – Trauma – Possible abruption – Hx anticoagulant use or bleeding disorder – Hx Female Genital Mutilation – Meconium in liquor – Placenta previa/ low placenta – Cervical cerclage (stitch in) – Diabetes'</p>

CPGs	The principal differences are
	<p>Instruction box 'Wait at least one minute post birth. Clamp cord at 10, 15 & 20 cm from baby – Cut cord between 15 and 20 cm clamps' replaces 'Wait at least one minute post birth then clamp cord at 10, 15 & 20 cm from baby – Cut cord between 15 and 20 cm clamps'</p> <p>Transport 'To Obstetric Unit'</p> <p><i>Medication updates</i></p> <p>Consider treatment 'Consider analgesia' replaces 'Consider Nitrous Oxide & Oxygen'</p>
<p>CPG 4/5/6.12.5 Umbilical Cord Complications</p>	<p><i>Deleted</i></p> <p>Mandatory sequence step 'Request Ambulance Control to contact GP/ midwife/medical team as required by local policy to come to scene or meet en route'</p> <p>Sequence step 'Attempt to slip the cord over the baby's head'</p> <p>Sequence step 'Ease the cord from around the neck as shoulders are delivered'</p> <p>Consider treatment 'Nifedipine 20 mg PO'</p> <p><i>Added</i></p> <p>Instruction box 'Use a hands off approach unless there are complications. Avoid touching cord. Avoid manipulation, traction and stimulation until baby is fully delivered'</p> <p>Instruction box 'Pre-alert hospital at earlier opportunity. Emergency caesarean section may be required for cord prolapse' replaces 'For prolapsed cord pre-alert hospital as emergency caesarean section will be required'</p> <p>Sequence step 'Avoid excessing manipulation and traction on the cord'</p> <p>Sequence step 'Apply additional clamps to cord on either side of the rupture' replaces 'Apply additional clamps to cord'</p> <p>Mandatory sequence step 'Mother to adopt head down in left lateral position (hips higher than head)' replaces 'Mother to adopt head down left lateral position'</p> <p>Sequence step 'Hold presenting part off the cord using fingers, rotate fingers as required' replaces 'Hold presenting part off the cord using fingers' and is a Paramedic and AP level skill</p> <p>Sequence step 'Minimal handling of cord and cover with sterile pad' replaces 'Maintain cord temperature and moisture'</p> <p>Consider treatment option 'If prolonged transport time (> 15 min) consider inserting an indwelling catheter into the bladder and run 500 mL of NaCl into the bladder and clamp catheter' replaces 'Consider inserting an indwelling catheter into the bladder and run 500 mL of NaCl into the bladder and clamp catheter' is a non-core element for AP level</p> <p>Mandatory sequence step 'Rapid transfer to Obstetrics Unit'</p> <p>Transport 'To Obstetric Unit'</p>

CPGs	The principal differences are
<p>CPG 4/5/6.12.6 Post Pregnancy Care (Including miscarriage and abortion)</p>	<p>The CPG is retitled 'Post Pregnancy Care (Including miscarriage and abortion)' (previously Postpartum Haemorrhage) The CPG entry point is updated to '≤ 6 weeks Post-partum' The CPG treatment pathway is significantly reorganised</p> <p><i>Deleted</i></p> <p>Instruction box 'Estimate blood loss' Instruction box 'Check/ask mother re multiple births prior to administration of Oxytocin' Sequence step 'Apply absorbent pad to perineum area' Sequence step 'Elevate lower limbs' Consider treatment 'Consider inserting a urinary catheter'</p> <p><i>Added</i></p> <p>Special instruction box 'If possibility of on-going pregnancy go to pregnancy CPG' Clinical finding 'PV Bleeding' Consider treatment 'Consider retained parts of conception as cause' Decision process 'Signs of shock' replaces 'Mother is haemodynamically unstable' Sequence step 'Uterine massage' replaces mandatory sequence step 'External massage of the uterus' and is introduced as an EMT level skill Consider treatment 'Consider breast feeding (If no contraindications)' Decision process 'Signs of sepsis' Transport 'To Obstetric Unit' Clinical finding 'Sepsis' Instruction box 'Additional sepsis symptoms – Low back pain – PV bleed – PV discharge' Clinical finding 'Delivery ≥ 20 weeks with; Headache, Visual disturbance, Dyspnoea, Oedema or seizure' Sequence step 'Measure BP' Decision process 'Eclamptic seizure or pre-eclampsia suspected (BP > 140/90 x 2)' 'Request ALS' Instruction box 'Suspect pre-eclampsia if above symptoms present and 2 elevated BP readings 15 min apart' Transport 'To General ED' Clinical finding 'Altered Mood' Consider 'Mental Health CPG' Sequence step 'Assess home environment & supports (report at handover)'</p> <p><i>Medication Updates</i></p> <p>Oxygen therapy is deleted</p>

CPGs	The principal differences are
CPG 4/5/6.13.1 Primary Survey Medical – Paediatric	<i>Added</i> Special instruction box 'Report findings as per Children First Act 2015 to ED staff and Tusla in a confidential manner' replaces 'Report findings as per Children First guidelines to ED staff and line manager in a confidential manner'
CPG 4/5/6.13.2 Primary Survey Trauma – Paediatric	<i>Added</i> Special instruction box 'Report findings as per Children First Act 2015 to ED staff and Tusla in a confidential manner' replaces 'Report findings as per Children First guidelines to ED staff and line manager in a confidential manner' Sequence step 'Jaw thrust' replaces sequence step 'Jaw thrust (Head tilt/ chin lift)'
CPG 4/5/6.13.4 Secondary Survey – Paediatric	<i>Added</i> Special instruction box 'Report findings as per Children First Act 2015 to ED staff and Tusla in a confidential manner' replaces 'Report findings as per Children First guidelines to ED staff and line manager in a confidential manner'
CPG 4/5.13.5 Foreign Body Airway Obstruction – Paediatric	The CPG treatment pathway is reorganised with the introduction of a cyclical process <i>Deleted</i> Sequence step '1 to 5 back blows followed by 1 to thrusts as indicated' Instruction box 'After each cycle of CPR open mouth and look for object. If visible attempt once to remove it' <i>Added</i> Cyclical process '1 to 5 back blows' - '1 to 5 thrusts (child - abdominal thrusts – infant – chest thrusts) - 'If patient becomes unresponsive' Decision process 'Effective' Sequence step 'Open mouth and look for object – If visible make one attempt to remove it' replaces sequence step 'Open mouth and look for object – If visible one attempt to remove it'
CPG 4/5/6.13.7 Abnormal Work of Breathing – Paediatric	The CPG is retitled Abnormal Work of Breathing – Paediatric (previously Inadequate Ventilations – Paediatric)

CPGs	The principal differences are
<p>CPG 4/5/6.13.9 Stridor – Paediatric</p>	<p><i>Added</i></p> <p>Consider treatment option ‘Consider humidified O₂ – as high a concentration as tolerated’ is a non-core element for EMT, paramedic and AP level and replaces sequence step ‘Humidified O₂ – as high a concentration as tolerated’.</p> <p>Mandatory sequence step ‘Do not insert anything into the mouth (other than PO medications for croup)’ replaces ‘Do not insert anything into the mouth’</p> <p>Instruction box ‘Signs of Croup may include; Hypoxia/ cyanosis – Marked respiratory distress – Stridor – Irritability or lethargy – Marked increased respiratory rate – If persistent treat as severe croup – If symptoms are intermittent treat as moderate croup’</p> <p>Instruction box ‘Maximum Dexamethasone administered within the past 72 hours not to exceed 600 mcg/kg’</p> <p>Medication Updates</p> <p>Oxygen therapy and sequence step ‘Humidified O₂ – as high a concentration as tolerated’ have been uncoupled</p>
<p>CPG 4/5/6.13.11 Glycaemic Emergency – Paediatric</p>	<p><i>Added</i></p> <p>Consider treatment option ‘Consider Ketone measurement’ is a non-core element for EMT, Paramedic and AP level</p> <p>Medication Updates</p> <p>Glucagon age specific doses IM are revised</p>
<p>CPG 4/5/6.13.13 Pain Management – Paediatric</p>	<p><i>Added</i></p> <p>Instruction box ‘Following Fentanyl IN the next dose may be either Fentanyl IN or Morphine IV’ replaces ‘Following Fentanyl IN the next dose may be either Fentanyl IV or Morphine IV but not both’.</p> <p>Instruction box ‘Morphine PO for ≥ 1 year olds only – Repeat Morphine at not < 2 min intervals prn to Max of 100 mcg/kg IV.’ replaces ‘Morphine PO for ≥ 1 year olds only – Repeat Morphine at not < 2 min intervals prn to Max of 0.1 mg/kg IV</p> <p>Instruction box ‘Repeat Ketamine PRN at not < 10 minutes.’ replaces ‘Repeat Ketamine once only at < 10 minutes prn.</p> <p>Instruction box ‘Poly-opiate administration should be avoided where possible – where multiple opiates are administered continuous patient monitoring is essential’</p> <p>Medication Updates</p> <p>Paracetamol PO dose is revised to 15 mg/kg</p>

CPGs	The principal differences are
CPG 4/5/6.13.14 Burns – Paediatric	<i>Added</i> Instruction box 'Should cool for another 20 minutes during packaging and transfer – Caution with hypothermia' replaces Instruction box 'Should cool for another 10 minutes during packaging and transfer – Caution with hypothermia'
CPG 4/5/6.13.16 External Haemorrhage – Paediatric	<i>Deleted</i> Paramedic skill flag from mandatory sequence step 'Apply tourniquet if limb injury' Paramedic skill flag from sequence step 'Depress proximal pressure point' Paramedic skill flag from sequence step 'Apply tourniquet' 'apply a tourniquet and/or' from EMT-BTEC Special Authorisation box <i>Added</i> Mandatory sequence step 'Apply and mark tourniquet if limb injury' replaces 'Apply tourniquet if limb injury' and is introduced as an EMT level skill Consider treatment option 'Consider wound closure clips for temporary closure if serious haemorrhage' is a non-core element for Paramedic and AP level Consider treatment option 'consider applying a dressing impregnated with haemostatic agent' is a Paramedic level skill Consider treatment option 'consider applying a dressing impregnated with haemostatic agent' is a non-core element for EMT level
4/5/6.13.19 Pyrexia – Paediatric	Medication Updates Paracetamol PO dose is revised to 15 mg/kg
CPG 4/5/6.13.20 Sepsis – Paediatric	The CPG is retitled Sepsis – Paediatric (previously Septic Shock – Paediatric) EMT level is added to this CPG The CPG treatment pathway is significantly reorganised The CPG entry point is updated to 'Patient generally unwell with suspected infection Temperature < 36°C or > 38.5°C' <i>Deleted</i> Sequence step 'Signs of Systemic Inflammatory Response Syndrome (SIRS)' Sequence step 'Could this be a severe infection?' Instruction box 'Normal ranges (ICTS)' Instruction box 'Give three' Instruction box 'If history of penicillin allergy assess the severity of the reaction and if not life-threatening, i.e. rash, proceed with Ceftriaxone' Instruction box 'If meningitis suspected ensure appropriate PPE is worn; Mask and goggles' Instruction box 'Signs of inadequate perfusion'

CPGs	The principal differences are
CPG 4/5/6.13.20 Sepsis – Paediatric (contd.)	<p><i>Added</i></p> <p>Sequence step ‘SpO₂, BP, RR, ETCO₂ & ECG monitoring’ replaces ‘ECG, SpO₂ & BP monitoring’</p> <p>Mandatory sequence step ‘Abnormal physiology? Source of infection considered’</p> <p>Decision process ‘Sepsis Red or Amber Flag +/- risk factors’</p> <p>Decision process ‘Evidence of inadequate perfusion’</p> <p>Instruction box ‘Titrate SpO₂ ≥ 94%’</p> <p>Instruction box ‘Sepsis Red Flag (≥ 1 item) – Altered mental status (P or U on AVPU) – Inappropriate tachycardia – Prolonged central capillary refill – Non-blanching rash – Hypotension – Clinical deterioration’</p> <p>Instruction box ‘Sepsis Amber Flag (≥ 1 item) – Inappropriate tachypnoea – Altered functional status – Practitioner concern – Parental concern – Vital signs deterioration – Risk factor(s) +/- Immunocompromised – Age ≤3 months – Chronic disease – Recent surgery – Break in skin (including chicken pox) – Indwelling line/device – Signs of infection in wound – Incomplete vaccination record’</p> <p>Sequence step ‘Monitor clinical condition; re-evaluate for possible sepsis if clinically indicated’</p> <p>Decision process ‘Clinical status improving’</p> <p>Decision process ‘Consider 2nd fluid bolus’</p> <p>Instruction box ‘High Consequence Infectious Disease (HCID) ensure appropriate PPE is worn; Long sleeve gown, Facemask, Eye protection’</p> <p>Special instruction box ‘If infection advise Triage nurse’ replaces ‘If SIRS + infection advise Triage nurse’</p> <p>Medication Update</p> <p>Paracetamol PO dose is revised to 15 mg/kg</p>
CPG 4/5/6.13.21 Allergic Reaction/ Anaphylaxis - Paediatric	<p>Medication Updates</p> <p>Chlorphenamine age specific doses IM are revised</p> <p>Adrenaline (1:1000) age specific doses IM are revised</p>
CPG 4.13.26 Post-Resuscitation Care – Paediatric	<p><i>Deleted</i></p> <p>Instruction box ‘Titrate O₂ to 96% - 98%’</p> <p><i>Added</i></p> <p>Sequence step ‘Avoid warming’ replaces sequence step ‘Prevent warming’</p>
CPG 4/5/6.14.2 VF or pVT – Adult	<p><i>Added</i></p> <p>Mandatory sequence step ‘Defibrillate’ replaces ‘Defibrillate (escalating energy)’</p> <p>Consider treatment option ‘Consider mechanical CPR assist’ replaces consider treatment ‘Consider mechanical CPR assist’ and is a non-core element for EMT, Paramedic and AP level</p>

CPGs	The principal differences are
CPG 4.14.3 Asystole – Adult	<i>Added</i> Consider treatment option 'Consider mechanical CPR assist' replaces consider treatment 'Consider mechanical CPR assist' and is a non-core element for EMT
CPG 4/5/6.14.5 Pulseless Electrical Activity – Adult	<i>Added</i> Consider treatment option 'Consider mechanical CPR assist' replaces consider treatment 'Consider mechanical CPR assist' and is a non-core element for EMT, Paramedic and AP level
CPG 4.14.6 Post-Resuscitation Care - Adult	<i>Medication Update</i> Consider treatment 'If Opioid OD suspected consider Naloxone dose and route specific to clinical level'
CPG 4.15.1 End of Life - DNAR	The CPG is retitled 'End of Life – DNAR' (previously End of Life – DNR)
CPG 4/5/6.16.3 Triage Sieve	<i>Deleted</i> Decision process 'Can casualty walk' Decision process 'Is casualty breathing' Mandatory sequence step 'Open airway one attempt' Decision process 'Breathing now' <i>Added</i> Decision process 'Catastrophic haemorrhage' Mandatory sequence step 'Apply and mark a tourniquet' Consider treatment option 'consider applying a dressing impregnated with haemostatic agent' is a non-core element for EMT level Decision process 'Is the casualty injured' Destination 'Survivor Reception Centre' Decision process 'Can the patient walk' Decision process 'Airway (open) & Breathing' Decision process 'Respond to Voice (AVPU)' Mandatory sequence step 'Recovery position'



Pre-Hospital
Emergency Care
Council



Published by Pre-Hospital Emergency Care Council

2nd Floor, Beech House, Millennium Park, Osberstown, Naas, Co. Kildare W91 TK7N

☎ 045 882042 ✉ info@phecc.ie

www.phecc.ie